



INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C09B 57/00, 23/00, G01N 33/58		A1	(11) International Publication Number: WO 97/40104
			(43) International Publication Date: 30 October 1997 (30.10.97)
(21) International Application Number: PCT/GB97/01105		(74) Agent: PENNANT, Pyers; Stevens Hewlett & Perkins, 1 Serjeants' Inn, Fleet Street, London EC4Y 1LL (GB).	
(22) International Filing Date: 21 April 1997 (21.04.97)			
(30) Priority Data: 96302783.4 19 April 1996 (19.04.96) EP		(81) Designated States: CA, JP, RU, US, European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).	
(34) Countries for which the regional or international application was filed: GB et al.		Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.	
(71) Applicant (for all designated States except US): AMERSHAM INTERNATIONAL PLC (GB/GB); Amersham Place, Little Chalfont, Buckinghamshire HP7 9NA (GB).			
(72) Inventors; and			
(75) Inventors/Applicants (for US only): HAMILTON, Alan, Lewis (GB/GB); "Ashdown", 15 Church Grove, Little Chalfont, Amersham, Buckinghamshire HP6 6SH (GB). WEST, Richard, Martin (GB/GB); 38 Pages Lane, Uxbridge, Middlesex UX8 1XT (GB). BRIGGS, Mark, Samuel, Jonathan (GB/GB); 109 Park Lane, Amersham, Buckinghamshire HP6 6NQ (GB). CUMMINS, William, Jonathan (GB/GB); 5 Thorntree Drive, Tring, Herts HP23 4BE (GB). BRUCE, Ian, Edward (GB/IE); 37 Greenare, Dunslaughlin, County Meath (IE).			
(54) Title: SQUARATE DYES AND THEIR USE IN FLUORESCENT SEQUENCING METHOD			
(57) Abstract Novel squarate dyes are described, and adducts of these dyes with biologically significant chemical species such as nucleosides or nucleotides. The adducts have useful properties for fluorescent nucleic acid sequencing methods.			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

SQUARATE DYES AND THEIR USE IN FLUORESCENT SEQUENCING METHOD

5 This invention concerns the use of a class of dyes in various biological applications. Some of the dyes are claimed as new compounds *per se*.

Background to the Invention

10 The development of automated fluorescent methods has led to increased data generation in DNA sequencing projects. Smith *et al* US 5,171,534 have described a fluorescent DNA sequencing system. Waggoner *et al* US 5,268,486 have described the properties of some conjugates of cyanine dyes and Middendorf US 5,230,781 and Patonay
15 EP 670 374 have described the use of various cyanine dyes in DNA sequencing. Berger *et al* European patent 214 847 has described the use of other cyanine dyes some of which contain squarate groups in assays which involve a specific binding partner. Other squarate dyes are described by Pease *et al* in USP 4,830,786 and subsequent divisional
20 patents, and by A J G Mank *et al* in Anal. Chem. 1995, 67, 1742-8. Cushman *et al* WO 93/09172 and Krutak *et al* WO94/19387 have described cyanine dyes containing squarate groups for use in thermoplastics and inks.

25 There is a need for methods of detecting biologically significant chemical species (hereafter biological molecules) at increased convenience and sensitivity in general and particularly for DNA sequencing and DNA mapping experiments.

Summary of Present Invention




In one aspect the invention provides an adduct of any biological molecule with a squarate dye as defined below. Examples of biological molecules are peptides, proteins, antibodies, polysaccharides and drugs. Preferably the biological molecule is a nucleoside or nucleotide or analogue or oligonucleotide. An analogue of a nucleoside or nucleotide may be a nucleoside or nucleotide derivative or other sugar-heterocycle which inhibits or mimics biological activity of normal nucleosides or nucleotides towards nucleic acid modifying or polymerising enzymes. An example of a nucleotide analogue is a chain terminator, such as a dideoxynucleotide, as used in sequencing reactions. The adduct may have the formula Q-N-CO-Sq, where Q is a biological molecule such as a nucleoside or nucleotide or analogue or oligonucleotide residue, and Sq is a residue of a squarate dye, the two being joined by an amide linkage formed between an amine group of Q and a carboxylate group of Sq. Alternatively the linkage may be formed between a functional group such as carboxylic acid, or derivatives thereof, isothiocyanate, maleimide, iodoacetamide, or phosphoramidite, and a nucleophile such as an amine, thiol, hydroxy or other group, as known for other nucleotide or oligonucleotide adducts, whereby either the nucleophilic or electrophilic reactive grouping may be attached to the dye. Alternatively the dye containing O-alkyl or O-alkenyl or O-alkynyl on the central moiety but with no other reactive functional group on the rest of the molecule may react with an amine or thiol or alcohol group to form a covalent linkage to a biological molecule.




In a further aspect the invention provides for the labelling of species immobilised on solid supports. One example of this may be an immobilised labelled oligonucleotide monomer or a difunctional derivatised dye with one arm bound to a solid support which is used for automated DNA synthesis. In a later step the labelled oligonucleotide is

cleaved from the support. Another application may be the labelling of a suitably derivatised solid support which is used in a heterogeneous luminescence based assay. In this instance the label may or may not be cleaved from the support.

5 In another aspect, the invention provides an improved fluorescent sequencing method, which comprises using an adduct as defined.

Squarate dyes are described in EP 214 847, the disclosure of which is incorporated herein by reference. A major aspect of the
10 present invention is concerned with a family of squarate dyes that are particularly suitable for use in the adducts and the improved fluorescent sequencing methods defined above. According to this aspect, the invention provides a squarate dye of the formulae (I), (II), (IIa), (III), (IV) or (IVa).





where each Z is independently O or S or CR¹₂,

n = 1 - 3,

R¹ is lower alkyl (1 - 4 carbon chain),

each R² is independently selected from electron donating


5 and electron withdrawing groups such as halogen, alkoxy, primary secondary and tertiary amino, nitro, SO₃⁻, and -R³-X, or is a branched or straight chain of up to 30 carbon atoms incorporating one to five positively charged nitrogen atoms,

each R³ is independently selected from: alkylene,

10 alkenylene and alkynylene (1 - 20 carbon chain), or is a branched or straight chain of up to 30 carbon atoms incorporating one to five ether oxygen atoms or arylene rings or positively charged nitrogen atoms,

at least one X is a nucleophilic functional group, such as OH, SH or NH₂, or alternatively a grouping capable of reacting with a
15 nucleophile, in which case X is preferably selected from the following:

CO₂H, activated carboxyl such as acid halide or anhydride, CO active ester, NCS, O phosphoramidite, NC(O)CH₂I and



20

any other X present is independently selected from H and SO₃⁻, and the residue of a squarate dye (whereby dimers and oligomers of the dyes shown as monomers of formula (I), (II), (IIa), (III), (IV) and (IVa) are envisaged) or other fluorochrome,

25

each of s and m is 0, 1 or 2,

A is O, NR⁴ or S,

R⁴ is alkyl, alkenyl, alkynyl or H, and

each Y is independently X or H.

At least one group R^2 may be SO_3^- in the compounds of formula (I), (II) and (IIa). Except when X is phosphoramidite (when SO_3^- groups are optional), preferably 1 to 5 SO_3^- groups are present to provide improved solubility in aqueous solvents.

The presence of sulphonic acid groups within the dye confers several advantages, namely increased water solubility, increased photostability, brightness and the potential reduction of interactions with surroundings.

Biological molecules such as proteins, antibodies, DNA and RNA are intrinsically water soluble to enable them to carry out their functions in a biological environment. There are well known procedures for isolating them by the addition of organic solvents, such as ethanol, to precipitate them from solution. To enable them to be labelled by reactive dyes, the dye molecules themselves must be either soluble in an aqueous environment or a mixed aqueous organic environment that does not precipitate or denature the biological molecule being labelled. The presence of sulphonic acid groups on the described squarate dyes greatly enhances their water solubility. Thus the squarate dyes synthesised for comparative studies in Example 13 lacking the sulphonic acids were found to require the addition of dioxan for labelling a H_2N -DNA primer as described in Example 5. Although a degree of successful labelling of DNA could be achieved by such procedures the relatively high organic content would likely cause either precipitation or denaturing of a protein. The squarate dye (13c) in Example 13 derived from benzindole derivatives was initially found to be non-fluorescent. The fluorescent properties were restored by boiling the dye in a 1% SDS detergent solution to prevent aggregation and hence self-quenching of fluorescence. Such aggregation is well known for cyanine dyes in the photographic industry (West, W and Pierce S. J. Phys Chem 69, 1894

(1965), Sturmer, D.M, Spec Top in Heterocyclic Chemistry, 30 (1974)).
When the corresponding sulphonated dye (2h) in Example 2 was studied
no such problems were encountered.

5 The presence of sulphonate groups on the squarate dye confers to
the dye an overall net negative charge. This assists in reducing non-
specific hydrophobic interaction with biological molecules. DNA and RNA
are by nature negatively charged due to the phosphate backbone, thus a
negatively charged dye will be repelled by electrostatic interactions and
10 limit any labelling to the specific reaction of the attached reactive
groups/nucleophile. Even after labelling the presence of sulphonate
groups on the dye in the dye conjugate will assist in minimising
hydrophobic interactions with any plastic components encountered in the
manipulation of the squarate dye conjugate. In sequencing applications,
15 particularly capillary gel sequencing, the negative charge on the dye will
prevent any adverse interaction with the capillary wall coating within the
capillary that would greatly distort the results, as the capillary wall coating
has already been optimised for the negatively charged DNA.

20 The presence of a sulphonic acid group on the aryl portion of the
dye imparts greater photostability to the dye. This is illustrated by the
tables in Example 13, which has determined the $t_{1/2}$ of the dyes when
exposed to a bright light source. All those squarate dyes lacking an aryl
sulphonate have the lowest $t_{1/2}$. The presence of two sulphonic groups
25 increases further the photostability. For comparison a commercially
available cyanine dye (Cy5™) has also been included in the tables.

 When the reactive group is a phosphoramidite the absence of a
sulphonic acid group is preferred but not essential. It is well known that
30 DNA/RNA can be synthesised on a solid support by using

phosphoramidite nucleotide monomer building blocks. There is also a range of phosphoramidite labels that can be used to attach haptens and dyes within a growing DNA chain or at the terminal 5' or 3' end dependent upon the users requirements. S.L. Beaucage *et al* in Tetrahedron **49**,
5 1925, (1993). Common to all these approaches is a protection strategy to ensure all reactive nucleophilic hydroxyl or amino functionalities within any phosphoramidite are protected to prevent reaction with the phosphorus (III) reagent during phosphoramidite activation and addition. On a DNA synthesiser these phosphoramidite additions are normally
10 carried out in organic solvent, typically acetonitrile.

It has been found that alcohol derivatives of the squarate dyes of this invention can be readily converted to the required phosphoramidite derivative without recourse to protection of the hydroxyl species on the
15 central cyclobutenediylum-1,3-diolate ring to which the indoleninium etc. intermediates have been coupled. This is an unexpected result and contrasts with the elaborate synthesis required to produce a fluorescein dye phosphoramidite performed by P. Theisen *et al* Tetrahedron Lett. **33**, 5033, (1992). The absence of a sulphonic acid group in the squarate dye
20 phosphoramidites is desirable so that acetonitrile can still be used as the reaction solvent of choice on the DNA synthesisers. It has also be found that phosphoramidites of the squarate dyes can be synthesised even when a sulphonic acid group is present. Both types of derivatives have been used to label a DNA primer and subsequently generate sequence
25 information - see Example 11.

The synthesis of the squarate dyes can be carried out either in a one step procedure or by the reaction of squaric acid or its derivatives with first one chosen intermediate and isolation of the "half dye" type structure with subsequent reaction with the intermediate of choice to
30 provide the required unsymmetrical dye. Example 1 provides a range of

intermediates that can be used to prepare any specific dye of choice. Substituents (R^2) on the aromatic rings can lead to significant effects on the properties of the dyes, e.g. wavelength shifts and stability. (See The Chemistry of Synthetic Dyes, Venkataraman, Academic Press NY 1971, vol.4, chapter 5, part iic, pages 228-240, particularly Table 1 on page 230.). The intermediates also provide a range of X derivatives that can be chosen e.g. OH, NH_2 which can both be readily converted into phosphoramidite or iodoacetamide respectively and carboxylic acid groups for conversion to activated carboxylic derivatives upon the required strategy for coupling to a biology molecule. Thus nucleotides such as 5'- aminoallyl dUTP require a succinimidyl ester squarate derivative where as a cysteine residue on a protein requires a maleimide or iodoacetimide squarate dye derivative. Those skilled in the art of conjugation will realise the conjugation strategies above are illustrative and are not meant to be limiting.

The variations induced by the variants in the substituents on the aromatic rings (R^2), or an increase in the number of conjugated aromatic rings (e.g. benzindole instead of indole) generally provide relatively subtle changes of wavelength and stability. A more significant change in wavelength properties can be induced by varying n in the central ring of the dye. These dyes are based on the squaric acid ($n = 1$), croconic ($n = 2$), and rhodizonic ($n = 3$), collectively termed squarate dyes herein for brevity. The value of n determines the approximate maximum excitation wavelength : 570-690nm $n = 1$, 690-790nm $n = 2$, 790-890nm $n = 3$.

The synthesis of unsymmetrical squarate dyes with functional groups or reactive groups attached to linker arms either on the aromatic ring as an substituent or off the nitrogen in the heteroaromatic ring is an achievable process as demonstrated in the examples. Generally taking advantage of symmetry can improve the overall synthetic yields in any given process. The modification on the central cyclobutenediylum -1,3-

diol at ring where $n=1$ and the corresponding derivatives where $n=2$ or 3 can provide an overall more efficient process as well as a novel labelling position. The Examples 7-10 provide processes wherein the squarate dyes are first converted to an ether derivative and then subsequently
5 further reacted to provide the required functionality for attachment to a biological molecule. The modification of the initial ether derivatives is not necessarily required for attachment to biological molecules as the ether derivative itself will react with amine groups. Modification of the central ring is not the only strategy that can be employed to increase ease of
10 synthesis. The mono-protection of difunctional squarate dyes (e.g. two identical X groups present) as in Examples 11h and 11k can also provide mono-reactive derivatives. This approach also allows for deprotection of the second functional group for the subsequent reaction to a second biological molecule, stationary phase or dye as required.

15 The modification of the central ring has also been found to alter the fluorescent properties of the dyes. Thus, the replacement of the initial O methyl substituent with a R^4NMe group has been found to dramatically reduce fluorescence providing for a quencher type dye. When the substituent on the aromatic portion of the dye (R^2) is a nitro
20 group the same sort of affect can be achieved.

These squarate dyes can be used in fluorescence energy transfer (ET). This technology is mediated by a dipole - dipole coupling between chromophores that results in resonance transfer of excitation from an excited donor chromophore to an acceptor chromophore (Forster,
25 T (1965) in Modern Quantum Chemistry, Istanbul lectures, part III Ed. Sinanoglur, O (Academic, New York) pp 93-137).

Fluorescence ET is a useful spectroscopic phenomenon that is well known in biological analysis, (Stryer, L, Ann. Rev. Biochem., (1978) 47 819-846; Cardullo, R A, *et al* Proc. Natl. Acad. Sci USA, (1988) 85
30 8790-8794; Ozak H *et al*, Nucleic Acids Res., (1992) 20, 5205-5214;

Clegg R M *et al*, Biochemistry (1992) 31 4846-4856 and Proc. Natl. Acad. Sci. USA, (1993) 90 p2994-2998; Selvin P R, Proc. Natl. Acad. Sci USA, (1994) 91 10024-10028).

In one example, the donor dye, absorbs light at the
5 wavelength of, for example, the appropriate laser. The energy emitted from this donor dye is transferred to a second dye, the acceptor dye. This acceptor dye emits the energy as fluorescence at the normal wavelength at which the acceptor dye emits. For example, a system based upon
10 squarate dyes derived from two indolinium intermediates, as a donor absorbing at ca. 633 nm, and an acceptor derived from benzindolinium intermediates, absorbing at ca. 665 nm, can be envisaged.

This principle has been used in many biological assays included DNA sequencing and analysis (Jingyue Ju *et al* Proc. Natl. Acad. Sci USA (1995) 92 p4347-4351).

15 In a further example the acceptor molecule may be chosen such that it quenches the energy emitted from the donor. The acceptor is then called a quencher. Such principles have been used in homogeneous gene detection assays. (Tyagi S *et al* Nature Biotechnology (1996) 14 (3) p303-308). Squarate dyes can be designed
20 which can be used as donors and/or acceptors or quenchers as described above.

Thus the invention also provides a fluorescent labelling complex comprising:

- a first or donor fluorochrome having first absorption and
25 emission spectra;
- a second or acceptor fluorochrome having second absorption and emission spectra, the wavelength of the emission maximum of said second fluorochrome being longer than the wavelength of the emission maximum of said first fluorochrome, and a portion of the
30 absorption spectrum of said second fluorochrome overlapping a portion of

the emission spectrum of said first fluorochrome;

- at least one linker for covalently attaching said first and second fluorochromes for transfer of resonance energy transfer between said first and second fluorochromes;

5 • a target bonding group capable of forming a covalent bond with a target compound;

wherein at least one of the said first and second fluorochromes is a squarate dye.

As demonstrated in the experimental section below,

10 squarate oligonucleotides conjugates can be used successfully in automated fluorescent DNA sequencing. In this application they can offer several potential advantages over other dyes which absorb at shorter and longer wavelengths:

1. The 632 nm red HeNe laser is significantly cheaper than the
15 Argon ion, GaAlAs, YAG and 594 nm HeNe lasers used in other DNA sequencers.

2. The optics and filters are much simpler and cheaper than diffraction grating, fibre optic and scanning confocal microscope arrangements used by other sequencers.

20 3. The longer excitation wavelength makes the use of soda lime glass plates possible, avoiding more expensive low-fluorescence borosilicate glass.

4. The red laser causes less background fluorescence from gel and buffer components which in turn increases signal to noise levels and
25 improves sensitivity.

5. The squarate dyes are more photostable than other dyes such as cyanines.

6. By the careful selection of dyes with differing spectral characteristics DNA could be sequenced within one track on a slab or
30 capillary gel based sequencing instrument.

7. By the careful manipulation of the overall dye charge versus the degree of dye lipophilicity it is possible to synthesis either pairs or sets of dyes such that they have a matched gel mobility shift upon conjugation to DNA. This provides for the ease of user analysis of all the raw

5 sequence data and reduces the reliance upon complex deconvolution algorithms and computer generated results.

The squarate dye may include a branched or straight chain of up to 30 atoms incorporating 1-5 positively charged nitrogen atoms. Preferably each positively charged nitrogen atom is provided by a
10 quaternary ammonium group, an imidazole group or a pyridinium group.

The above illustrates how the squarate dyes can be modified by the addition of charged residues, for example the sulphonate grouping which provides for negatively charged dyes. In certain applications neutral or positively charged dyes are either necessary or
15 advantageous. The addition of various numbers of quaternary nitrogen species to the dyes will provide overall positively charged dyes if no sulphonate groups are present or overall negatively, neutral or positively charged dyes if sulphonate groups are present. The above combined with the ability to vary the wavelength of any given dye by the choice of
20 dye starting materials and/or squaric acid derivatives allows for the synthesis of matched dyes suitable for 2D gel applications as outlined in Waggoner *et al.* WO 96/33406.

EXPERIMENTAL EXAMPLES**EXAMPLE 1****Synthesis of intermediates for dye synthesis**

The starting quaternised indolenines and related derivatives were prepared according to the methods of R.B.Mujumdar *et al.*

Bioconjugate Chemistry, 1993, 4, 105, G. Patonay *et al.* J.Org. Chem. 1995, 60, 2391 and E. Barni *et al.* Heterocyclic Chem, 1985, 22, 1727. A

representative example is included in each case.

Potassium 2,3,3-trimethylindolenine-5-sulphonate (1a)

To a 2l three necked round bottomed flask equipped with a mechanical stirrer was added acetic acid (300 ml), 3-methyl-2-butanone (168 ml, 1.6 mol) and 4-hydrazinobenzene sulphonic acid (100 g, 0.53 mol). This was then heated under reflux for 3h and then cooled, with stirring, overnight. The resulting pink precipitate was collected by filtration and then dried *in vacuo* at 60°C.

The crude product was converted to the title potassium salt by dissolution in methanol followed by addition of a saturated solution of KOH in *iso*-propyl alcohol. The precipitated yellow solid was collected by filtration and dried *in vacuo* at 60°C.

δ_H (270 MHz, D₂O) 7.15 (1H, s), 7.11 (1H, dd, $J = 7.0, 1.2$ Hz), 6.52 (1H, d, $J = 7.0$ Hz), 2.21 (3H, s), 1.38 (6H, s).

Potassium 1-(4-sulphonatobutyl)-2,3,3-trimethylindoleninium-5-sulphonate (1b)

The potassium salt (1a) (11.0 g, 40 mmol) and 1,4-butane sultone (6.5 g, 48 mmol) were mixed together in 1,2-dichlorobenzene (50 ml) and then heated with stirring at 110°C for 8h. The mixture was then

cooled overnight. The excess liquid was decanted off and the residue triturated with iso-propyl alcohol, filtered and dried *in vacuo* at 60°C. This was then HPLC purified (C-18, H₂O/MeOH).

δ_H (270 MHz; D₂O) 8.08 (1H, s), 7.96 (1H, dd, $J = 9.0, 1.2$ Hz),
5 7.31 (1H, d, $J = 9$ Hz), 4.48 (2H, t, $J = 7.5$ Hz), 2.23 (2H, t, $J = 7.5$ Hz), 2.04 (3H, s), 1.95 (2H, m), 1.48 (6H, s), 1.35 (2H, m).

1-(5-Carboxypentyl)-2,3,3-trimethylindoleninium-5-sulphonate (1c)

Synthesised by an analogous method to (1b)

10 δ_H (270 MHz; D₂O) 8.10 (1H, s), 7.99 (1H, dd, $J = 9.0, 1.2$ Hz),
7.29 (1H, d, $J = 9.0$ Hz), 4.48 (2H, t, $J = 7.5$ Hz), 2.29 (2H, t, $J = 7.4$ Hz), 2.01 (2H, m), 1.61 (6H, s), 1.32 - 1.60 (4H, m); (270 MHz; DMSO d_6) 8.03 (1H, s),
7.96 (1H, d, $J = 8.24$ Hz), 7.84 (1H, d, $J = 8.24$ Hz), 4.46 (2H, t, $J =$
7.14Hz), 2.86 (3H, s), 2.23 (2H, t, $J = 7.14$ Hz), 1.84 (2H, m), 1.55 (6H, s),
15 1.41 (4H, m)

6-Bromo-3-oxahexanoic acid

To a stirred solution of glycolic acid (2.03 g, 26 mmol) and
1,3-dibromopropane (6.41 g, 32 mmol) in THF (50 ml) was added sodium
20 hydride (1.56 g, 65 mmol). This was stirred at room temperature for 16h.
The reaction mixture was quenched with dilute HCl (1.0 M, 100 ml) and
then extracted into chloroform (3x50 ml). This was then washed with brine,
dried, filtered and evaporated to dryness *in vacuo*. The residue was
purified by chromatography (SiO₂, CHCl₃/MeOH) to yield the title compound
25 (2.91 g, 57%).

δ_H (270 MHz; CDCl₃) 3.95 (2H, s), 3.41 (2H, t, $J = 7.0$ Hz), 3.25 (2H, t, $J = 7.0$ Hz), 1.57 (2H, m).

1-(5-Carboxy-4-oxapentyl)-2,3,3-trimethylindoleninium-5-sulphonate (1d)

Synthesised by an analogous method to (1b) using 6-bromo-3-oxahexanoic acid

5 δ_H (D₂O) 8.10 (1H, s), 7.99 (1H, dd, $J = 9.0, 1.2\text{Hz}$), 7.29 (1H, d, $J = 9.0\text{Hz}$), 4.48 (2H, t, $J = 7.5\text{Hz}$), 3.95 (2H, s), 3.25 (2H, t, $J = 7.0\text{Hz}$), 1.61(6H, s), 1.57 (2H, m).

1-Ethyl-2,3,3-trimethylindoleninium-5-sulphonate (1e)

10 Synthesised by an analogous method to (1b)

δ_H (270 MHz; DMSO d₆) 8.02 (1H, s), 7.94 (1H, d, $J = 8.24\text{Hz}$), 7.83 (1H, d, $J = 8.24\text{Hz}$), 4.48 (2H, q, $J = 7.14\text{Hz}$), 2.85(3H, s), 1.54 (6H, s), 1.44 (3H, t, $J = 7.14\text{Hz}$).

15 **1-Butyl-2,3,3-trimethylindoleninium-5-sulphonate (1f)**

Synthesised by an analogous method to (1b)

δ_H (270 MHz; D₂O) 0.85 (2H, t), 1.35 (2H, m), 1.50 (6H, s), 1.83 (2H, quin), 4.36 (2H, t), 7.80 (1H, m), 7.91 (1H, m) and 8.02(1H, app s).

20

1-Ethyl-2,3,3-trimethylindoleninium iodide (1g)

Synthesised by an analogous method to (1b)

25 δ_H (270 MHz; DMSO d₆) 7.97 (m, 1H), 7.84 (m, 1H), 7.63 (m, 2H), 4.49 (q, 2H, $J = 7.14\text{Hz}$), 2.85 (s, 3H), 1.54 (s, 6H), 1.45 (t, 3H, $J = 7.14\text{Hz}$).

1-(4-Sulphonatobutyl)-2,3,3-trimethylindoleninium-5-acetic acid (1h)

Synthesised by an analogous method to (1b)

δ_H (270 MHz; DMSO d_6) 1.52 (6H, s), 1.74 (2H, quin), 1.96 (2H, quin), 2.83 (3H, s), 3.44 (2H, br s), 3.74 (2H, s), 4.47 (2H, br t), 7.51 (1H, d), 7.70 (1H, s) and 7.96 (1H, d).

5 **1-(5-Carboxypentyl)-2,3,3-trimethylindoleninium bromide (1l)**

Synthesised by an analogous method to (1b)

δ_H (270 MHz; DMSO- d_6) 7.89 (m, 1H), 7.86 (m, 1H), 7.63 (m, 2H), 4.47 (t, 2H; $J = 7.42\text{Hz}$), 2.86 (s, 3H), 2.24 (t, 2H; $J = 7.14\text{Hz}$), 1.83 (m, 2H), 1.55 (s, 6H), 1.46 (m, 4H).

10

1-Butyl-2,3,3-trimethylindoleninium-5-acetic acid iodide (1j)

Synthesised by an analogous method to (1b)

δ_H (270 MHz; CD_3OD) 1.05 (3H, t), 1.45-1.60 (2H, m), 1.85-2.05 (2H, m), 3.75 (2H, s), 4.50 (2H, t), 7.57 (1H, d), 7.70 (1H, s) and 7.80 (1H, d).

15

1-Ethyl-2,3,3-trimethylbenzindoleninium iodide (1k)

Synthesised by an analogous method to (1b)

δ_H (270 MHz; CDCl_3) 1.52 (3H, t), 1.78 (12H, s), 2.97 (3H, s), 4.64 (2H, q), 7.75 (2H, quin) and 8.20-8.50 (4H, m).

20

1-(5-Carboxypentyl)-2,3,3-trimethylbenzindoleninium bromide (1l)

Synthesised by an analogous method to (1b)

δ_H (270 MHz; CDCl_3) 1.50-1.95 (4H, m), 1.85 (12H, s), 2.05 (2H, quin), 2.37 (2H, t), 4.65 (2H, t), 7.70 (1H, t), 7.80 (1H, t), 8.05 (1H, d), 8.17 (1H, d), 8.25 (1H, d) and 8.35 (1H, d).

25

5-Iodopentyl acetate

5-Chloropentyl acetate (26.5g, 0.16mol) was added to a solution of sodium iodide (45g, 0.30mol) in dry acetone (200ml). The

30

resulting pale yellow solution was heated at reflux for 65h, during which time a white solid separated (sodium chloride). The final mixture was cooled to room temperature and filtered to remove the NaCl, which was washed with acetone and diethyl ether, then dried under vacuum at 50°C.

5 Expected yield of NaCl = 9.4g; isolated yield = 9.47g (100%).

The filtrate and washings were combined and the solvent removed under reduced pressure; the residue was partitioned between water and diethyl ether. The ether layer was retained, washed with aqueous sodium thiosulphate solution and brine, then dried (Na₂SO₄),
10 filtered and the solvent removed under reduced pressure, to give *titled compound* as a yellow-tinged liquid, 39g (95%).

δ_H (270MHz, CDCl₃) 1.50 (2H, m), 1.65 (2H, m), 1.85 (2H, m), 2.05 (3H, s, CH₃-COO-), 3.18 (2H, t, I-CH₂-), 4.06 (2H, t, -CH₂-OAc).

15 **1-(5-Acetoxy-pentyl)-2,3,3-trimethylindoleninium iodide (1m)**

5-Iodopentyl acetate (3.84g, 15mmol) was added to freshly distilled 2,3,3-trimethylindolenine (2.39g, 15mmol); this mixture was then heated at 100°C for 4h under nitrogen atmosphere, to give a reddish gum. After cooling to room temperature this was triturated repeatedly with
20 diethyl ether and dried under high vacuum to give a viscous reddish gum, 6.1g (98%). This material was used directly, without further purification.

3-(5-Acetoxy-pentyl)-1,1,2-trimethylbenz(e)indoleninium iodide (1n)

5-Iodopentyl acetate (19.4g, 78mmol) was heated to 50°C,
25 then 1,1,2-trimethyl-1H-benz(e)indole (16.2g, 77mmol) was added. This mixture was heated at 100°C for 4.5h before cooling to room temperature. The solidified melt was dissolved in 10% methanol / dichloromethane (100ml), then diluted with diethyl ether (400ml) to precipitate the product. After 30mins stirring the pale green crystals were collected, washed with
30 ether and dried to give (1n), 33.1g (92%).

MS (MALDI-TOF): 339

δ_H (300MHz, $CDCl_3$) 1.55 (2H, m, $-CH_2-$), 1.68 (2H, m, $-CH_2-$), 1.83 (6H, s, indole CMe_2), 1.97 (3H, s, CH_3-COO-), 2.0 (2H, m, $-CH_2-$), 3.2 (3H, s, indole C_2-CH_3), 4.02 (2H, t, indole $N-CH_2-$), 4.78 (2H, t, $-CH_2-OAc$), 7.6-7.7 5 (2H, m), 7.82 (1H, d), 8.03 (2H, m), 8.08 (1H, d).

N-(3-Bromopropyl)triethylammonium bromide

1,3-Dibromopropane (20.0g, 100mmol) and triethylamine (5.06g, 50mmol) were mixed in dry toluene (50ml). This solution was heated at 100°C under nitrogen atmosphere for 4h, during which time a thick white solid precipitated. The mixture was then cooled and the solid collected by filtration, washed with toluene and ether and dried under vacuum at 50°C to give the *titled compound* 5.0g (36%).

δ_H (300MHz, DMSO) broad peaks. 1.17 (9H, $3 \times N^+-CH_2-CH_3$), 2.15 (2H, $BrCH_2CH_2CH_2-$), 3.26 (8H, $4 \times N^+-CH_2-$), 3.62 (2H, $Br-CH_2-$).

1-((3-Triethylammonium)propyl)-2,3,3-trimethylindolium dibromide (1o)

Freshly distilled 2,3,3-trimethylindolenine (0.8g, 5mmol) and N-(3-bromopropyl)-triethylammonium bromide (1.52g, 5mmol) were mixed and placed under an argon atmosphere. The mixture was then heated at 140°C for 1.5h, giving a deep red viscous melt, which solidified to a glass on cooling. It was ground to a powder under diethyl ether; this was collected by filtration, triturated with boiling acetone and recrystallised from methanol / acetonitrile to give the *title compound* as a pale pink powder, 795mg (34%).

δ_H (300MHz, DMSO) 1.22 (9H, t, J 6.6Hz, $3 \times N^+-CH_2-CH_3$), 1.55 (6H, s, indole C_3Me_2), 2.21 (2H, m, $-CH_2CH_2CH_2-$), 2.92 (3H, s,

indole C2-Me), 3.27 (6H, q, J 6.6Hz, $3 \times \text{N}^+-\text{CH}_2-\text{CH}_3$), 3.51 (2H, ~t, $-\text{CH}_2-\text{NEt}_3$), 4.57 (2H, ~t, indole N^+-CH_2-), 7.64 (2H, m), 7.86 (1H, d, J 6.5Hz), 8.12 (1H, d, J 7.3Hz).

5 **1-(3-Triethylammonium)propyl)-2,3,3-trimethylbenzothiazolium dibromide (1p)**

Synthesised by an analogous method to (1o)

Recrystallised from 1-butanol / acetone.

δ_{H} (300MHz, D_2O) 1.15 (9H, t, $3 \times \text{N}^+-\text{CH}_2-\text{CH}_3$), 2.3 (2H, m, -
10 $\text{CH}_2\text{CH}_2\text{CH}_2-$), 3.25 (6H, q, $3 \times \text{N}^+-\text{CH}_2-\text{CH}_3$), 3.45 (2H, ~t, $-\text{CH}_2-\text{NEt}_3$), 4.6 (2H, ~t, indole N^+-CH_2-), 7.4 (1H, t), 7.6 (1H, t), 8.0 (1H, d.), 8.1 (1H, d).

3-(3-Aminopropyl)-1,1,2-trimethylbenz(e)indolium bromide.HBr (1q)

Prepared according to Patonay *et al.*, *J. Org. Chem.*, (1995),

15 60, 2391.

δ_{H} (300MHz, DMSO) 1.77 (6H, s, indole C1Me_2), 2.24 (2H, m, $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$), 2.98 (3H, s, C2 CH_3), 3.10 (2H, app q, $-\text{CH}_2-\text{NH}_3^+$), 4.72 (2H, t, J 7.0Hz, indole $\text{N}-\text{CH}_2-$), 7.70-7.81 (2H, m) + 8.21-8.39 (4H, m) = $6 \times$ benzoindeole aryl-H, 8.03 (3H, broad s, $-\text{NH}_3^+$).

20

3-(3-N-Phthalimidopropyl)-1,1,2-trimethylbenz(e)indolium bromide (1r)

Synthesised by an analogous method to (1q)

δ_{H} (300MHz, CDCl_3) 1.84 (6H, s, indole C1Me_2), 2.44 (2H, m, -
25 $-\text{CH}_2-\text{CH}_2-\text{CH}_2-$), 3.24 (3H, s, C2 CH_3), 3.86 (2H, t, J 7.1Hz, $-\text{CH}_2-$ phthalimide), 4.99 (2H, t, J 7.2Hz, indole $\text{N}-\text{CH}_2-$), 7.58-7.76 (7H, m), 7.94-8.02 (3H, m).

N-Ethyl-2-methylbenzothiazolium iodide (1s)

Synthesised by an analogous method to (1b)

δ_H (270MHz, $CDCl_3$) 1.6 (6H, t, benzothiazole N- CH_2-CH_3),
3.5 (3H, s, benzothiazole C2- CH_3), 5.0 (2H, q, benzothiazole N- CH_2-
5 CH_3), 7.75 (1H, t) + 7.85 (1H, t) + 8.1 (1H, d) + 8.3 (1H, d) = 4x
benzothiazole aryl-H. δ_H (270 MHz; CD_3OD) 1.60 (3H, t, NCH_2CH_3), 3.23
(3H, s, Me), 4.86 (2H, q, NCH_2CH_3), 7.82 (1H, t, ArH), 7.95 (1H, t, ArH),
8.28 and 8.32 (each 1H, overlapping d, ArH).

10 **N-(5-Carboxypentyl)-2-methylbenzothiazolium bromide (1t)**

Synthesised by an analogous method to (1b)

δ_H (270 MHz; CD_3OD) 1.57, 1.71 and 2.00 (each 2H, m),
2.26 (2H, t, CH_2CO_2H), 4.77 (2H, t, NCH_2), 7.82 and 7.93 (each 1H, t,
ArH) and 8.21-8.40 (2H, m, ArH).

15

3-Ethyl-1,1,2-trimethylbenz(e)indoleninium-7-sulphonate (1u)

Synthesised by an analogous method to (1b)

δ_H (300 MHz; D_2O) 1.45 (3H, t, CH_2CH_3), 1.57 (6H, s, CMe_2),
4.46 (2H, q, CH_2CH_3), 7.79, 7.84, 8.04 and 8.18 (each 1H, d, ArH) and
20 8.26 (1H, d, ArH).

1-(5-Carboxypentyl)-1,1,2-trimethylbenz(e)indoleninium-7-sulphonate (1v)

Synthesised by an analogous method to (1b) to give insoluble
25 grey powder which was used directly with structural confirmation obtained
from product dyes.

3-Ethyl-6,8-disulphonato-1,1,2-trimethylbenz(e)indolium tosylate (1w)

Synthesised by an analogous method to (1b)

δ_H (300MHz, D₂O) 1.39 (3H, app t, indole N-CH₂-CH₃), 1.53 (6H, s, indole C2Me₂), 2.18 (3H, s, tosylate -CH₃), 4.40 (2H, q, *J* 7.5Hz, indole N-CH₂-CH₃), 7.15+ 7.50 (each 2H, \approx d, 4 \times tosylate aryl-H), 7.92 (1H, d, *J* 9.5Hz), 8.42 (1H, s), 8.49 (1H, s), 8.79 (1H, d, *J* 9.5Hz).

5

3-(5-Carboxypentyl)-6,8-disulphonato-1,1,2-trimethylbenz(e)indolium bromide (1x)

Synthesised by an analogous method to (1b)

δ_H (300MHz, D₂O) 1.30 (2H, m), 1.48 (2H, m), 1.59 (6H, s, indole C2Me₂), 1.83 (2H, m), 2.19 (2H, t, *J* 7.2Hz, -CH₂-CO₂H), 4.41 (2H, t, *J* 7.5Hz, indole N-CH₂-), 7.92 (1H, d, *J* 9.5Hz), 8.40 (1H, s), 8.53 (1H, s), 8.79 (1H, d, *J* 9.5Hz).


10

1-Benzyl-2,3,3-trimethylindoleninium-5-sulphonate (1y)


15

Synthesised by an analogous method to (1b)


δ_H (300MHz, CD₃OD) 1.66 (6H, s, CMe₂), 5.85 (2H, s, PhCH₂), 7.37-7.44 (5H, m, PhCH₂), 7.77 (each 1H, d, *J* 8.4, 4-CH), 7.91 (1H, dd, *J* 1.5 and 8.4, 6-CH) and 8.11 (1H, d, *J* 1.5, 7-CH).




- (1g) R=Et
 (1i) R=(CH₂)₅CO₂H
 (1m) R=(CH₂)₅OAc
 (1o) R=(CH₂)₃N⁺(Et)₃




- (1b) R=(CH₂)₄SO₃⁻
 (1c) R=(CH₂)₅CO₂H
 (1d) R=(CH₂)₃OCH₂CO₂H
 (1e) R=Et
 (1f) R=Bu
 (1y) R=Bn




- (1h) R=(CH₂)₄SO₃⁻
 (1j) R=Bu




- (1w) R=Et
 (1x) R=(CH₂)₅CO₂H



- (1k) R=Et
 (1l) R=(CH₂)₅CO₂H
 (1n) R=(CH₂)₅OAc
 (1q) R=(CH₂)₃NH₂·HBr
 (1r) R=3-phthalimidopropyl



- (1u) R=Et
 (1v) R=(CH₂)₅CO₂H



- (1p) R=(CH₂)₃N⁺(Et)₃
 (1s) R=Et
 (1t) R=(CH₂)₅CO₂H

EXAMPLE 2**Synthesis of sulphonic acid substituted squarate dyes from indoleninium intermediates**

5 For the invention dyes the exact nature of the counter-ions was not determined.

2-(1-(5-Carboxypentyl)-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-4-(1-(4-sulphonatobutyl)-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (2a)

A mixture of (1b) (0.56 g, 1.35 mmol) and squaric acid (0.48 g, 4.05 mmol) in 2-ethoxyethanol (10 ml) was heated under reflux for 20h. The reaction mixture was then evaporated to dryness *in vacuo* and purified by HPLC (C-18, H₂O/MeOH) to furnish an intermediate mono-adduct of
15 squaric acid ($\lambda_{\text{max}}(\text{H}_2\text{O}) = 425\text{nm}$). Said adduct (0.27 g, 0.65 mmol) and (1c) (0.19 g, 0.71 mmol) in 2-ethoxyethanol (10 ml) were heated under reflux for 16h. The 2-ethoxyethanol was removed *in vacuo* and the residue was purified by HPLC (C-18, H₂O/MeOH) to afford the title compound (2a).

δ_{H} (270 MHz; D₂O) 7.81 (2H, s), 7.65 (2H, d, $J = 7.5\text{Hz}$), 7.31-
20 7.43 (2H, m), 6.27 (2H, m), 4.11 (4H, m), 2.89 (2H, t, $J = 7.0\text{Hz}$), 2.45 (2H, t, $J = 7.0\text{Hz}$), 1.4-2.0 (22H, m); $\lambda_{\text{max}}(\text{H}_2\text{O}) = 631\text{nm}$.

2-(1-(5-Carboxypentyl)-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-4-(1-ethyl-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (2b)

Squaric acid (121 mg, 1.06 mmol) in n-butanol (50 ml) was heated at reflux until all the squaric acid had dissolved. Then (1c) (500 mg, 1.06 mmol) was added portionwise over 2h and the mixture was maintained at reflux for a further 2h. 1-Ethyl-2,3,3-trimethylindoleninium-
30 5-sulphonate (1e) (439 mg, 1.06 mmol) was added portionwise over 1h.

After heating at reflux for a further 24h the mixture was cooled to ambient temperature and concentrated *in vacuo*. Purification of the residue by HPLC (C18, H₂O/MeOH) afforded the *title compound* as its n-butyl ester.

λ_{\max} (MeOH) 636 nm; λ_{\min} (MeOH) 652 nm

5 δ_{H} (270 MHz; CD₃OD) 0.92 (3H, t, $J = 7.3$ Hz), 1.38-1.93 (13H, m), 1.78 (12H, s), 2.34 (2H, t, $J = 7.1$ Hz), 4.04 (2H, t, $J = 6.6$ Hz), 4.08-4.28 (4H, m), 6.02 (1H, s), 6.04 (1H, s), 7.25-7.35 (2H, m) and 7.82-7.92 (4H, m).

To a solution of the above dye butyl ester (13 mg) in water (2 ml) was added 1M KOH solution (0.5 ml). After stirring at ambient temperature overnight HPLC purification of the mixture (PRP-1, H₂O/MeOH) afforded the title compound as the potassium salt.

λ_{\max} (MeOH) 636 nm; λ_{\min} (MeOH) 647 nm

15 δ_{H} (270 MHz; CD₃OD) 1.40 (3H, t, $J = 7.1$ Hz), 1.45-1.58 (2H, m), 1.63-1.93 (4H, m), 1.78 (12H, s), 2.19 (2H, t, $J = 7.4$ Hz), 4.09-4.28 (4H, m), 6.03 (1H, s), 6.04 (1H, s), 7.25-7.36 (2H, m) and 7.85-7.92 (4H, m)

2-(1-(5-Carboxypentyl)-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-4-(1-butyl-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (2c)

The title compound was prepared as its butyl ester in a similar manner to (2b) butyl ester, *vide supra*, from (1c), squaric acid and (1f).

λ_{\max} (MeOH) 638 nm

25 δ_{H} (270 MHz; CD₃OD) 0.91 (3H, t, $J = 7.3$ Hz), 1.02 (3H, t, $J = 7.3$ Hz), 1.26-1.44 (2H, m), 1.44-1.64 (6H, m), 1.64-1.92 (6H, m), 1.77 (12H, s), 2.34 (2H, t, $J = 7.1$ Hz), 4.06 (2H, t, $J = 6.6$ Hz), 4.15 (4H, app br t, $J = 6.9$ Hz), 6.02 and 6.03 (each 1H, s), 7.28 (2H, app d, $J = 4.1$ Hz), 7.86 (2H, app d, $J = 4.1$ Hz) and 7.88 (2H, app s).

30 Subsequent hydrolysis gave the title compound as the

potassium salt which was used in later studies.

2-(1-(5-Carboxypentyl)-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-4-(1-ethyl-3,3-dimethyl-2-

indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (2d)

The title compound was prepared as its butyl ester in a similar manner to (2b) butyl ester, *vide supra*, from (1c), squaric acid and (1g).

λ_{max} (MeOH) 630 nm

δ_{H} (270 MHz; CD₃OD) 0.92 (3H, t), 1.27-1.92 (13H, m), 1.76 (12H, s), 2.34 (2H, t), 4.04 (2H, t), 4.11 (2H, br t), 4.23 (2H, br q), 5.95 and 6.05 (each 1H, s), 7.19-7.32 (3H, m), 7.33-7.45 (1H, m), 7.49 (1H, d) and 7.80 to 7.88 (2H, m).

Saponification of the above butyl ester (ca. 11mg) and subsequent purification by chromatography (C18, H₂O/MeOH), in a similar manner to (2b), *vide supra*, afforded the title compound as the potassium carboxylate (8 mg)

δ_{H} (270 MHz; CD₃OD) 1.42 (3H, t), 1.46-1.58 (2H, m), 1.62-1.91 (4H, m), 1.76 and 1.78 (each 6H, s), 2.19 (2H, t), 4.11 (2H, br t), 4.21 (2H, br t), 5.96 and 6.04 (each 1H, s), 7.20-7.34 (3H, m), 7.34-7.45 (1H, m), 7.45-7.51 (1H, m) and (7.83-7.89 (2H, m).

2-(1-(4-Sulphonatobutyl)-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-4-(1-(4-sulphonatobutyl)-3,3-dimethyl-5-carboxymethyl-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (2e)

Squaric acid (55 mg, 0.484 mmol) was dissolved in a mixture of acetic acid (5 ml), pyridine (5 ml) and acetic anhydride (500 μ l) to give a clear yellow-orange solution. (1b) (200 mg, 0.484 mmol) was added in two portions (over 15 min) to give a green solution. After a

further 15 min 1-(4-sulphonatobutyl)-2,3,3-trimethylindoleninium-5-acetic acid (1h) (85 mg) was added followed by a further portion (85 mg) after 35 min. The resultant dark green solution was stirred at ambient temperature for ca. 20h, concentrated *in vacuo* and redissolved in H₂O/MeOH.

- 5 Purification of the crude dye by HPLC (C18, H₂O/MeOH), with isolation of the middle one of three major blue components, afforded the *title compound* as the pyridinium salts. Percolation through an acid exchange resin (Dowex 50W, ca. 10 ml), eluting with water (25 ml), and concentration *in vacuo* of the eluent afforded the title compound as the
10 free acids.

λ_{max} (MeOH) 638 nm

δ_{H} (270 MHz; D₂O) 1.35 and 1.45 (each 6H, s), 1.75 (8H, br s), 2.80 (4H, app br t), 3.40 (2H, br s), 3.95 (4H, br m), 7.00-7.30 (4H, m) and 7.70 (2H, br s)

15

2-(1-Ethyl-3,3-dimethyl-5-sulphonato-2-indolinyldenemethyl)-4-(1-(5-carboxypentyl)-3,3-dimethyl-2-indolinyldenemethyl)-cyclobutenediylum-1,3-diolate (2f)

- The title compound was prepared as its butyl ester in a
20 similar manner to (2b) butyl ester, *vide supra*, from (1e), squaric acid and (1i).

- δ_{H} (270 MHz; CD₃OD) 0.92 (3H, t, CH₂CH₂CH₃), 1.00-1.90 (13H, m), 1.82 (12H, s, 2×CMe₂), 2.33 (2H, t, CH₂CO₂H), 4.02 (2H, t, CO₂CH₂), 4.08-4.25 (4H, m, 2×CH₂N), 5.96 and 6.03 (each 1H, s, vinylH),
25 7.16-7.55 (5H, series m, ArH) and 7.82-7.94 (2H, m, ArH).

Subsequent hydrolysis afforded the free acid.

- δ_{H} (270 MHz; CD₃OD) 1.38 (3H, t, CH₂CH₃), 1.55 (2H, m), 1.62-1.95 (4H, m), 1.76 (12H, s, 2×CMe₂), 2.20 (2H, t, CH₂CO₂H), 4.10-4.25 (4H, m, 2×NCH₂), 5.95 and 6.05 (each 1H, s, vinylH), 7.18-7.50 (5H, series m, ArH) and 7.80-7.89 (2H, m, ArH).
30

2-(1-Benzyl-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-4-(1-(5-carboxypentyl)-5-sulphonato-3,3-dimethyl-2-indolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (2g)

The title compound was prepared as its butyl ester in a similar manner to (2b) butyl ester, *vide supra*, from (1y), squaric acid and (1c).

δ_H (270 MHz; D₂O) 0.39 (3H, t, CH₂CH₂CH₃), 0.70-1.50 (22H, m), 1.88 (2H, br t, CH₂CO₂H), 3.58 (2H, br t, CO₂CH₂), 3.84 (2H, br, NCH₂CH₂), 4.96 (2H, br, PhCH₂), 5.63 and 5.67 (each 1H, overlapping br s, vinylH), 6.70-7.20 and 7.56-7.78 (11H, m, ArH).

Subsequent hydrolysis afforded the free acid.

δ_H (270 MHz; CD₃OD) 1.52 (2H, m), 1.63-1.90 (4H, m), 1.75 and 1.82 (each 6H, s, 2×CMe₂), 4.16 (2H, br t, NCH₂CH₂), 5.37 (2H, br s, PhCH₂), 6.03 and 6.07 (each 1H, s, vinylH), 7.16-7.40 (7H, m, ArH) and 7.72-7.93 (4H, m, ArH).

2-(1-(5-Carboxypentyl)-3,3-dimethyl-5-sulphonato-2-benzindolinyliidenemethyl)-4-(1-ethyl-3,3-dimethyl-5-sulphonato-2-benzindolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (2h)

The title compound was prepared in a similar manner to (2e) from (1u), (1v) and squaric acid in a two step procedure with isolation of the intermediate 'half dye'.

λ_{max} (MeOH) = 664 nm

MALDI-TOF (C₄₂H₄₃N₂S₂O₈ requires M⁺ 802) 824 (M⁺+Na), 711


2-(1-(5-Carboxypentyl)-3,3-dimethyl-5-sulphonato-2-benzindolinylidenemethyl)-4-(1-ethyl-3,3-dimethyl-2-benzindolinylidenemethyl)-cyclobutenediylum-1,3-diolate (2i)

The title compound was prepared in a similar manner to (2b) from (1k), (1v) and squaric acid.

λ_{max} (MeOH)=662 nm

δ_{H} (270 MHz; CD₃OD) 1.15 (3H, t, CH₂CH₃), 1.25 (2H, m), 1.44 (2H, m), 1.88 (2H, m), 1.99 (12H, s, 2×CMe₂), 2.25 (2H, br t, CH₂CO₂H), 4.19 (2H, br t, NCH₂CH₂), 4.27 (2H, br q, CH₂CH₃), 5.99 and 6.04 (each 1H, s, vinylH), 7.41 (1H, app t, ArH), 7.56 (3H, m, ArH), 7.92 (2H, app t, ArH), 8.01 (2H, app t, ArH), 8.24 (2H, m, ArH) and 8.41 (1H, s, ArH).

For examples of dyes bearing free hydroxyl groups see Examples 11 and 12.



(2a) $\text{R}_1 = (\text{CH}_2)_4\text{SO}_3^-$

$\text{R}_2 = (\text{CH}_2)_5\text{CO}_2\text{H}$

(2b) $\text{R}_1 = (\text{CH}_2)_5\text{CO}_2\text{H}$


$\text{R}_2 = \text{Et}$

(2c) $\text{R}_1 = (\text{CH}_2)_5\text{CO}_2\text{H}$

$\text{R}_2 = \text{Bu}$

(2g) $\text{R}_1 = \text{Bn}$

$\text{R}_2 = (\text{CH}_2)_5\text{CO}_2\text{H}$




(2d) $\text{R}_1 = (\text{CH}_2)_5\text{CO}_2\text{H}$

$\text{R}_2 = \text{Et}$


(2f) $\text{R}_1 = \text{Et}$

$\text{R}_2 = (\text{CH}_2)_5\text{CO}_2\text{H}$



(2e) $\text{R}_1 = \text{Bu}$

$\text{R}_2 = (\text{CH}_2)_4\text{SO}_3^-$



(2h) $R_1 = (CH_2)_5CO_2H$

$X = SO_3^-$

(2i) $R_1 = (CH_2)_5CO_2H$

$X = H$

EXAMPLE 3

5 Synthesis of sulphonic acid substituted squarate dyes from indoleninium and thiazolinium intermediates

10 2-(1-(5-Carboxypentyl)-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-4-(3-ethyl-2-benzothiazolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (3a)

Synthesised in an analogous manner to Example (2b) from intermediates (1c), (1s), squaric acid and additional quinoline as base to give a crude mixture of dye products. Hydrolysis of the crude material, purification by HPLC and percolation through H^+ exchange resin afforded the title compound.

$\lambda_{max}(\text{MeOH})$ 636 nm

$\delta_H(270 \text{ MHz; DMSO-}d_6)$ 1.18-1.82 (9H, m), 1.64 (6H, s, 2×Me), 3.96 (2H, br), 4.45 (2H, br), 5.57 and 6.09 (each 1H, br s, vinyl-CH), 7.11 (1H, m, ArH), 7.40 (1H, m, ArH), 7.48-7.70 (3H, m, ArH), 7.77 (1H, m, ArH) and 8.07 (1H, d, ArH)


2-(3-Ethyl-2-benzothiazolinyliidenemethyl)-4-(1-(4-sulphonatobutyl)-3,3-dimethyl-5-carboxymethyl-2-indolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (3b)

Synthesis in an analogous method to Example (3a) from intermediates


- 5 (1h) and (1s), with purification by HPLC only, afforded the mono-quinolinium salt of the *title compound*.

λ_{\max} (MeOH) 642nm

δ_H (270 MHz; CD₃OD) 1.43 (3H, t, CH₂CH₃), 1.68 (6H, s, 2×Me), 1.96 (4H, br), 2.91 (2H, br, CH₂SO₃⁻), 3.63 (2H, s, ArCH₂), 4.05
 10 (2H, br, NCH₂CH₂), 4.40 (2H, br q, NCH₂CH₃), 5.75 and 6.07 (each 1H, br s, vinyl-CH), 7.13 (1H, d, ArH), 7.23 (1H, d, ArH), 7.32 (1H, s, ArH), 7.38 (1H, t, ArH), 7.46-7.66 (2H, m, ArH), 7.75-7.95 (3H, m, ArH), 8.04 (1H, t, ArH), 8.18 (2H, t, ArH), 8.86 (1H, d, ArH) and 9.05 (1H, d, ArH)



(3a) R₁=(CH₂)₅CO₂H
 R₂=Et



(3b) R₁=(CH₂)₄SO₃⁻
 R₂=Et

EXAMPLE 4**Synthesis of the succinimidyl esters of the carboxylic acid derivatives of squarate dyes****A representative procedure for activation is as below, which is given for example (2b)**

To a solution of carboxylic acid derivative (2b) (potassium salt) (8 mg, 0.01 mmol) in DMF (1 ml) was added diisopropylethylamine (10.5 μ l, 0.06 mmol) and O-(N-succinimidyl)-N,N,N',N'-tetramethyluronium tetrafluoroborate (TSTU) (9 mg, 0.03 mmol). The reaction was monitored by HPLC (C18, 25 mM pH7 Phosphate buffer/MeCN) and upon completion was sealed under nitrogen and stored in a freezer until required for labelling experiments.

The follow squaric acid dyes were activated in an analogous manner:

(2a, c-h), (3a) and (3b).

EXAMPLE 5**Use of squaric acid derived fluorophores for automated fluorescent DNA sequencing****Primer synthesis**

An M13 (-20) universal 18mer primer was prepared according to established methods on an ABI model 394 DNA synthesiser. The 5' terminus of the oligonucleotide was modified by coupling an aliphatic amine group in the final synthesis cycle by means of a trifluoroacetamide protected-aminolinker amidite [Pharmacia Biotech].

The primer sequence used was NH₂-tgtaaaacgaacggccagt. The crude

primer was deprotected in 30% NH_4OH for 16 hours at 55°C then purified free of organic by-products and amines using phenol/chloroform extraction followed by ethanol precipitation. DNA was dissolved in water at 10D unit/ μl and stored at -70°C.

5

Primer labelling.

N-hydroxy succinimidyl esters of the squarate dyes were supplied in DMF or MeCN at 4 - 10 mg/ml. To label the primers, 40 μl of dye solution was mixed with 50 μl of 100mM sodium phosphate buffer at pH 8.0 and 10 μl [100D or 300 μg] of NH_2 -oligonucleotide. For squaric acid dyes not bearing any sulphonic acid groups (therefore with an overall neutral charge) and which are less soluble in 40% DMF, 1,4 dioxan was added to 10 - 20% v/v final concentration in order to keep the dye in solution and raise the labelling efficiency. Reactions were allowed to continue at +4°C in the dark for between 6 - 16h. Reactions were stopped by addition of sodium acetate pH 5.0 to a final concentration of 300 mM followed by 3 aqueous volumes of 99.8% ethanol to precipitate the oligonucleotide and remove unincorporated dye. After centrifugation at 13000 g for 15 minutes the DNA pellet was washed in 80% ethanol, dried *in vacuo* and dissolved in 100 μl of TE buffer at pH 8.0.

Primer purification

Squaric acid dye labelled primers were separated from residual, non-covalently attached dye and from unlabelled primer by HPLC on a 24 cm x 0.5 cm Spherisorb ODS2 column with a 5 μm support size. A 1 ml/minute gradient from 95% solvent A / 5% solvent B to 30% solvent A / 70% solvent B was used. A = 0.1M sodium acetate pH6.8, B = acetonitrile. Fractions absorbing at both 260 nm and 635 nm were collected and dried in a vacuum centrifuge. Fractions were pooled in 100 μl of TE buffer and ethanol precipitated as above to desalt them prior

to sequencing.

DNA sequencing

2 pmol of squarate dye-primer was mixed with 0.2 pmol of
5 M13 mp8 phage DNA in a volume of 25 μ l. The primer / template mixture
was divided into four 6 μ l aliquots designated A, C, G & T. 2 μ l of the
appropriate enzyme nucleotide / buffer / premix from Amersham kit
RPN2437 [Thermo Sequenase™ labelled primer sequencing kit] was
added and the reactions were placed in Perkin-Elmer GeneAmp PCR
10 system 9600 thermocycler. Samples were taken through 25 cycles at
95°C / 30seconds : 60°C / 30seconds. After thermocycling, 3 μ l of loading
dye containing 90% formamide / 5mM EDTA was added, the reactions
were concentrated in vacuum centrifuge for 10 minutes and then heat
denatured at 80°C for 2 minutes before loading on 6.1M urea / 5%
15 HydroLink gel in a Vistra DNA sequencer 725.

Electrophoresis and detection

A Vistra DNA sequencer 725 was modified by replacement
of the laser with a 30 mW Helium-Neon 632 nm tube and HV power
20 supply. The optical filter was changed to a 645 nm long-pass. No other
alterations were necessary to detect SQ5 fluorescence. Gels were run for
8 - 10 hours at 1400 V and maintained at or above 35°C throughout.
Image data was analysed using Vistra V2.01 software and Molecular
Dynamics' ImageQuaNT program.

25

Results

All of the Examples (2a-h) and (3a-b) gave sequence data.
The signal strength and peak resolution values were comparable to a
commercially available cyanine dye (Cy5™) commonly used for
30 sequencing. Base-calling accuracies of more than 98% to 550 bases are

typical.

EXAMPLE 6

5 The use of dye labelled oligonucleotides in two colour detection

An 18mer oligonucleotide was labelled separately with two cyanine dyes (6a) and (6b) and an invention dye (2f). The excitation and emission maxima of these dye oligonucleotide conjugates are shown in Table 1. 100 fmoles of each labelled oligonucleotide was loaded in the
10 following pairs into the same well onto a 19% denaturing acrylamide gel:

1. (6a) and (6b)
2. (6a) with (2f)
3. (2f) with (6b)

The labelled oligonucleotides were also loaded individually.

15 After electrophoresis under denaturing conditions the gel was scanned in a prototype scanning fluorescence instrument with a 633nm helium neon laser excitation light source. Fluorescent emission was collected in 3 sequential scans. The first scan with a 645nm RG filter (Schott) in place collected all the fluorescent light produced. The second
20 and third scans were with 660nm df₃₀ and 700nm eflp filters (Omega Optical) to collect and discriminate the fluorescence from each dye.

Each pair of dyes was identified from the 645nm RG filter image of the gel and sampled in a fluorochrome separation algorithm to identify the spectral properties of each dye. The images from each scan
25 were processed to produce overlaid 2 colour images.

These images show that (6a) with (6b) and (2f) with (6b) can be used as dye 2 colour pairs. These dyes are efficiently excited at the 633nm excitation wavelength and can be spectrally separated and identified for 2 colour applications.

Notes


The 645nm RG is a red glass long pass filter which excludes light below 645nm.

The 660 df₅₀ filter is a band pass filter centred at 660nm with
5 +/- 15 nm tolerance *i.e.* collects a band of light from 645-675nm.


The 700 eflp filter is a long pass filter which should exclude light below 700nm *i.e.* collects from 700nm upwards.

Table 1**10 Spectral Properties of Dye-Oligonucleotide Conjugates**

DYE 18 MER.	EX _{max} IN TE ₈ BUFFER	EM _{max} IN TE ₈ BUFFER
2f	630 nm	642 nm
6a	650 nm	660 nm
6b	680-685 nm	700 nm



(6a) $R_1 = (CH_2)_4SO_3^-$
 $R_2 = (CH_2)_3OCH_2CO_2H$



(6b) $R_1 = (CH_2)_4SO_3^-$
 $R_2 = (CH_2)_3OCH_2CO_2H$

EXAMPLE 7

5 Modifications of the central cyclobutenediylum-1,3-diolate ring of squarate dyes

2,4-Bis(1-ethyl-3,3-dimethyl-2-indolinyldenemethyl)-cyclobutenediylum-1,3-diolate (7a)

10 1-Ethyl-2,3,3-trimethylindoleninium iodide (**1g**) (3.15g, 10mmol), squaric acid (0.57g, 5mmol) and n-butanol (40ml) were mixed and heated at reflux for 16h. The solvent was then evaporated and the product dye (**7a**) isolated by flash chromatography (0-5% MeOH / CH₂Cl₂). Yield = 1.7g (75%)

15 λ_{max} (MeOH) = 628nm, λ_{ex} = 626nm, λ_{em} (MeOH) = 635nm

δ_H (270MHz, $CDCl_3$) 7.4-6.9 (8H, m), 6.0 (2H, s), 4.1 (4H, q), 1.8 (12H, s), 1.4 (6H, t).

Also isolated was the half-dye species 3-but xy-4-(1-ethyl-3,3-dimethyl-2-indolinyliidenemethyl)-cyclobut-3-en-1,2-dione (7b).

5 λ_{max} (MeOH) = 424nm.

δ_H (270MHz, $CDCl_3$) 7.4-6.8 (4H, m), 5.4 (1H, s), 4.85 (2H, t), 3.9 (2H, q), 1.9 (2H, m), 1.7 (6H, s), 1.5 (2H, m), 1.35 (3H, t), 1.0 (3H, t).

10 **3-Methoxy-2,4-bis(1-ethyl-3,3-dimethyl-2-indolinyliidenemethyl)-cyclobutenediylum-1-olate methosulphate (7c)**

The squarylium dye (7a) (1.0g, 2.2mmol) was dissolved in chloroform (20ml) to give a deep blue solution. To this was added dimethyl sulphate (5ml); the mixture was then heated at reflux for 16h.
15 The resulting solution was cooled, washed well with water, then dried (Na_2SO_4), filtered and evaporated to low volume. Diethyl ether was then added slowly, with scratching, up to about 100ml. This caused the methylated product (7c) to crystallize out as metallic green needles, which were collected by filtration, washed with fresh ether and dried under
20 vacuum. Yield = 1.24g (95%).

λ_{max} (MeOH) = 630nm, λ_{ex} (625), λ_{em} (MeOH)=638nm

δ_H (270MHz, $CDCl_3$) 7.43-7.38 (4H, m), 7.30 (2H, d), 7.18 (2H, d), 4.86 (3H, s), 4.3 (4H, q), 3.74 (3H, s), 1.73 (12H, s), 1.46 (6H, t).

(7c) is reactive, the O-Me group being readily replaced by an
25 alcohol or an amine. This is illustrated below.

3-Butoxy-2,4-bis(1-ethyl-3,3-dimethyl-2-indolinyliidenemethyl)-cyclobutenediylum-1-olate (7d)

The dye (7c) (50mg, 0.086mmol) was dissolved in n-butanol
30 (3ml) and heated at 70°C, with monitoring by t.l.c. (silica, 15% MeOH /

CH₂Cl₂). Once all the starting dye had been consumed (3-4h) the solution was cooled and evaporated. The residue was dissolved in chloroform (1ml) and diluted with diethyl ether (15ml); after standing in the freezer for 16h the product (7d) separated as metallic green needles.

5 Yield = 36mg (anion undetermined).

λ_{max} (MeOH) = 628nm.

δ_{H} (270MHz, CDCl₃) 7.5-7.0 (8H, m), 5.9 (2H, s), 5.2 (2H, t), 4.3 (4H, q), 4.1 (2H, t), 2.1 (2H, m), 1.8 (12H, s), 1.75-1.4 (m), 1.1 (3H, t), 0.9 (3H, t).

10

3-Butylamino-2,4-bis(1-ethyl-3,3-dimethyl-2-indolinyliidenemethyl)-cyclobutenediylum-1-olate (7e)

The dye (7c) (200mg, 0.35mmol) was dissolved in dichloromethane (20ml). To the stirred solution was added a solution of n-butylamine in dichloromethane (approx. 1 drop per ml CH₂Cl₂), in 0.1ml
15 portions. After each addition, the mixture was analyzed by t.l.c. (silica, 15% MeOH / CH₂Cl₂), until no more starting dye was present. The mixture was then evaporated and the residue purified by flash chromatography (silica, 4-10% MeOH / CH₂Cl₂). This gave 154mg of the
20 *title dye (7e)* as an amorphous powder after evaporation (anion undetermined).


λ_{max} (MeOH) = 646nm.

δ_{H} (270MHz, CDCl₃) 9.5 (1H, broad t), 7.6-6.9 (8H, m), 6.5 (1H, s), 5.7 (1H, s), 4.7 (2H, q), 4.0 (2H, q), 3.8 (2H, broad q), 1.9 (2H, m),
25 1.8 (12H, 2x s), 1.6 (2H, m), 1.5 (3H, t), 1.4 (3H, t), 1.0 (3H, t).

3-(7-(4,4'-Dimethoxytrityloxy)-6-hydroxy-4-oxaheptyl)amino-2,4-bis(1-ethyl-3,3-dimethyl-2-indolinyldenemethyl)-cyclobutenediylum-1-olate (7f)

The dye (7c) (58mg, 0.1mmol) was dissolved in
5 dichloromethane (4ml); to this was added a solution of 7-(4,4'-
dimethoxytrityloxy)-6-hydroxy-4-oxaheptylamine (EP 0527184B1)
(ca.0.3mmol in 2ml in dichloromethane) in portions, until t.l.c. analysis
(silica, 15% MeOH / CH₂Cl₂) indicated complete reaction. The solution
was then evaporated and the residue purified by flash chromatography
10 (silica, 4-10% MeOH / CH₂Cl₂). Evaporation gave 85mg of the product
(7f) as a violet-blue foam (anion undetermined).

λ_{\max} (CH₂Cl₂) = 652nm; λ_{\max} (CH₂Cl₂+CCl₃CO₂H) = 652
+504nm.




(7a) $R_1=R_2=Et$ $R_3=O^-$

(7c) $R_1=R_2=Et$ $R_3=OMe$ $MeOSO_3^-$

(7d) $R_1=R_2=Et$ $R_3=OBu$

(7e) $R_1=R_2=Et$ $R_3=NHBu$



(7f)

The above were used in DNA sequencing experiments
 5 using methods similar to those outlined in **Example 5**. The results were excellent. However, the method of labelling was not via the succinimidyl ester but by direct displacement of the dye ether linkage by the 5' amino substituent on the primer.

EXAMPLE 8**An active ester derivative of a cyclobutenediylum-1,3-diolate ring modified squarate dye****3-(N-Methyl-N-(3-carboxypropyl))amino-2,4-bis(1-ethyl-3,3-dimethyl-2-indolinyldenemethyl)-cyclobutenediylum-1-olate chloride (8a)**

4-Methylaminobutanoic acid.HCl (154mg, 1.0mmol) was mixed with methanol (10ml). To this was added a solution of tetrabutylammonium hydroxide in methanol (1.0M, 2.0ml, 2.0mmol). The methylated dye (7c) was then added (289mg, 0.5mmol) and the mixture stirred for a further 15min. This solution was poured into chloroform and washed three times with a large volume of water, then with saturated aqueous sodium bicarbonate solution, and finally with 0.1M HCl. The organic solution was dried (Na₂SO₄), filtered and evaporated; the residue was then purified by flash chromatography (silica, 5-20% MeOH / CH₂Cl₂), to give 212mg (72%) of the *title dye* (8a) as a violet-blue powder.

λ_{\max} (MeOH) = 660nm, λ_{ex} = 657nm, λ_{em} (MeOH) = 670nm

δ_{H} (270MHz, CDCl₃) 7.5-7.0 (8H, m), 6.0 (2H, s), 4.3 (4H, q), 3.85 (3H, s), 2.6 (2H, m), 2.2 (2H, m), 1.7 (12H, s), 1.4 (6H, s).

EXAMPLE NO.	VISIBLE ABSORBANCE (AU)	RELATIVE FLUORESCENCE INTENSITY
(7a)	0.489	713
(8a)	0.486	21


Table 8.1 Example of decrease in fluorescence for dye (8a)

Synthesis of the succinimidyl ester of the dye (8a)

The dye (8a) (59mg, 0.1mmol) was dissolved in dichloromethane (2ml). To this solution was added a solution of N-hydroxysuccinimide (12mg, 0.1mmol) in acetonitrile (2ml), followed by a solution of N,N'-dicyclohexylcarbodiimide in dichloromethane (1.0M, 0.12ml, 0.12mmol). This mixture was stirred for 24h, then it was filtered and evaporated. The residue was purified by flash chromatography (silica, 5-20% MeOH / CH₂Cl₂) to give the *active ester* as a solid with a metallic red lustre. Yield = 55mg (80%).

λ_{max} (MeOH) = 660nm

δ_{H} (270MHz, CDCl₃) 7.6-7.0 (8H, m), 6.0 (2H, s), 4.3 (4H, »q), 4.0 (2H, broad s), 3.7 (3H, s), 2.9 (2H, m), 2.7 (4H, s), 2.3 (2H, m), 1.7 (12H, s), 1.4 (6H, app t).



(8a)

EXAMPLE 9**Modifications to the central cyclobutenediylum-1,3-diolate ring of a thiazolinium based squarate dye**

5

2,4-Bis-(3-ethyl-2-benzothiazolinyldenemethyl)-cyclobutenediylum-1,3-diolate (9a)

3-Ethyl-2-methylbenzothiazolium iodide (**1s**) (1.55g, 5.0mmol), squaric acid (0.285g, 2.5mmol), quinoline (1.5ml) and n-butanol (20ml) were mixed and heated at reflux for 7h under nitrogen atmosphere. The resulting dark green mixture was cooled in the fridge overnight, then the solid collected by filtration. This solid was washed with a little ice-cold methanol, then diethyl ether, and dried under vacuum to give the *title dye* (**9a**) as a dark powder with a metallic olive-green lustre. Yield = 0.73g (68%).

λ_{max} (MeOH) = 650nm. λ_{em} (MeOH) 660nm.

δ_{H} (270MHz, CDCl_3) 7.6-7.0 (8H, m), 5.9 (2H, s), 4.2 (4H, q), 1.4 (6H, t).

20 **3-Methoxy-2,4-bis(3-ethyl-2-benzothiazolinyldenemethyl)-cyclobutenediylum-1-olate methosulphate (9b)**

Dye (**9a**) (0.73g, 0.7mmol) was dissolved in chloroform (30ml); to the resultant deep blue solution was added dimethyl sulphate (3ml). This mixture was heated at reflux for 7h, then left to stand for three days. It was then washed with water; the organic layer was retained and the aqueous layer extracted with more chloroform. The combined organic extracts were dried (MgSO_4), filtered and the solvent removed under reduced pressure, to a volume of ca. 15ml. Diethyl ether (40ml) was added to precipitate the product, which was collected by filtration, washed with ether and dried under vacuum to give the *titled dye* (**9b**) (1.0g, 100%).

T.l.c. (silica; 15% methanol / dichloromethane. (9a), $R_f = 0.75 \rightarrow$ (9b), $R_f = 0.3$).

λ_{\max} (MeOH) 632nm; λ_{ex} (MeOH) 649nm, λ_{em} (MeOH) 658nm.

5 δ_H (300MHz, CDCl_3) 1.29 (6H, t, J 7.0Hz, 2x benzothiazole $\text{N-CH}_2\text{-CH}_3$), 3.36 (3H, s, methosulphate -CH_3), 4.44 (4H, q, J 7.0Hz, 2x benzothiazole $\text{N-CH}_2\text{-CH}_3$), 4.51 (3H, s, squarate -OMe), 6.00 (2H, s, 2x methine $=\text{CH-}$), 7.37 (2H, app t), 7.53 (2H, app t), 7.72 (2H, d, J 8.4Hz), 8.01 (2H, d, J 8.1Hz).

10

3-Butylamino-2,4-bis(3-ethyl-2-benzothiazolinyliidenemethyl)-cyclobutenediylum-1-olate methosulphate (9c)

Methylated dye (9b) (100mg) was mixed with dichloromethane (30ml) and n-butylamine (0.2ml). This mixture was
15 stirred at room temperature for 30mins, during which time all the solid dissolved. T.l.c. (silica; 15% methanol / dichloromethane. (9b), $R_f = 0.3 \rightarrow$ (9c), $R_f = 0.45$). The solvent was removed under reduced pressure and the blue product isolated by flash chromatography (silica; 4-10% methanol / chloroform); the crude product gum was triturated with diethyl ether to
20 give a solid with a metallic red lustre, 43mg.

λ_{\max} (MeOH) 658nm; λ_{ex} (MeOH) 657nm, λ_{em} (MeOH) 671nm.

δ_H (300MHz, CDCl_3) 0.96 (3H, t, J 7.3Hz, butyl -CH_3), 1.36-1.51 (8H, m, 2x benzothiazole $\text{N-CH}_2\text{-CH}_3$ + $\text{-CH}_2\text{-}$), 1.83 (2H, m, $\text{-CH}_2\text{-}$),
25 3.56 (2H, q, J 7.1Hz, $\text{-HN-CH}_2\text{-}$), 4.11+4.47 (each 2H, q, J 7.0Hz, 2x benzothiazole $\text{N-CH}_2\text{-CH}_3$), 5.631 (1H, s, methine $=\text{CH-}$), 7.08-7.59 (9H, m, 8x benzothiazole aryl-H + methine $=\text{CH-}$), 10.17 (1H, broad app. t).

3-(N-methyl-N-(3-carboxypropyl))amino-2,4-bis(3-ethyl-2-benzothiazolinyliidenemethyl)-cyclobutenediylum-1-olate chloride (9d)

4-(Methylamino)butanoic acid.HCl (0.31g, 2.0mmol) was dissolved in methanol (10ml). To this solution was added a solution of tetra-n-butylammonium hydroxide in methanol (1M, 4ml, 4.0mmol), followed by dye (9b) (0.58g, 1.0mmol). The deep blue solution that resulted was stirred for 40mins; t.l.c. (silica; 15% methanol / dichloromethane. (9b), $R_f = 0.3 \rightarrow$ (9d), $R_f < 0.1$).

The solvent was removed under reduced pressure and the residue triturated with water. The metallic bronze solid was collected by filtration, washed with water and acetone, then dried under vacuum to give the acid dye (9d), 0.435g (82%).

λ_{max} (MeOH) 670nm; λ_{ex} (MeOH) 670nm, λ_{em} (MeOH) 686nm.

δ_H (300MHz, CD_3OD) 1.31 (6H, t, 2x benzothiazole $N-CH_2-CH_3$), 1.90 (2H, m, $-CH_2-CH_2-CH_2-$), 2.23 (2H, app t, $-CH_2-CO_2H$), 3.30 (3H, s, $N-CH_3$), 3.45 (2H, app t, $MeN-CH_2-$), 4.28 (4H, broad s, 2x benzothiazole $N-CH_2-CH_3$), 5.80 (2H, s, 2x methine $=CH-$), 7.23 (2H, d), 7.36 (4H, broad m), 7.67 (2H, m).

EXAMPLE NO.	VISIBLE ABSORBANCE (AU)	RELATIVE FLUORESCENCE INTENSITY
(9a)	0.490	1140
(9d)	0.485	124

Table 9.1 Example of the decrease in fluorescence for dye (9d)

3-(3-(*tert*-Butoxycarbonylamino)propylamino)-2,4-bis(3-ethyl-2-benzothiazolinyldenemethyl)-cyclobutenediylum-1-olate methosulphate (9e)


Synthesised in an analogous manner to (9c) using *tert*-butyl-
5 *N*-(3-aminopropyl)carbamate.

λ_{max} (MeOH)=656 nm

δ_{H} (300MHz, CDCl₃) 1.40 (9H, s, CMe₃), 1.48 (6H, t, J 6.6, 2×CH₂CH₃), 1.97 (2H, m, NHCH₂CH₂), 3.30 (2H, br q, CH₂NH), 3.67 (2H, br q, CH₂NH), 3.74 (3H, s, MeOSO₃⁻), 4.13 and 4.44 (each 2H, br q, CH₂CH₃), 5.69 (1H, s, vinylH), 5.89 (1H, br t, NHBOC), 6.58 (1H, s, vinylH), 7.12-7.60 (8H, series m, ArH) and 8.83 (1H, br t, vinylNH).

DNA sequencing experiments

Dye primer synthesis and subsequent sequencing
15 experiments were carried out in a similar manner to that outlined in Example 5 with labelling by either the succinimidyl ester or the ether derivative of the above dyes. The results were excellent.



- (9a) R=O⁻
 (9b) R=OMe MeOSO₃⁻
 (9c) R=NHBU
 (9d) R=N(Me)CH₂CH₂CH₂CO₂H Cl⁻
 (9e) R=NH(CH₂)₃NHBOC

EXAMPLE 10**Synthesis of a centrally modified croconic acid based dye****5 Croconic acid dye (10a)**

The dye (10a) was prepared according to the procedure outlined in US Patent 3793313 (1974).

λ_{max} (MeOH) 754nm

10 Methylation of (10a) to give (10b)

Dye (10a) (250mg, 0.54mmol) was dissolved in chloroform (12ml) to give a dark olive-green solution. To this was added dimethyl sulphate (1.5ml) and the mixture was heated at 60°C for 4h, giving a purple solution. T.l.c. (silica; 15% methanol / dichloromethane. (10a) yellow-green, dries to blue, $R_f = 0.6 \rightarrow$ (10b) purple, $R_f = 0.4$). The
15 solvent was removed under reduced pressure to a volume of ca. 2ml, then it was diluted with diethyl ether. The precipitated solid was collected by filtration, washed with more ether and dried under vacuum to give (10b), 330mg (100%).

20 λ_{max} (MeOH) 764nm

Coupling of (10b) to n-butylamine to give (10c)

Methylated dye (10b) (20mg) was dissolved in dichloromethane (5ml); to this was added n-butylamine (1 drop). The
25 colour of the solution changed quickly from purple to an orange-brown. T.l.c. (silica; 15% methanol / dichloromethane. (10b) purple, $R_f = 0.4 \rightarrow$ (10c) orange-brown, dries to purple, $R_f = 0.55$). The solution was purified by flash chromatography (silica; 10% methanol / dichloromethane) to give the amino dye (10c), 15mg.


λ_{\max} (MeOH) 768nm

δ_H (270MHz; CDCl₃) 0.98 (3H, t, CH₃CH₂CH₂), 1.50-2.00 (10H, m, 2×CH₂CH₃ and NCH₂CH₂CH₂), 4.30 (2H, m, NHCH₂), 4.50 and 4.90 (each 2H, br q, 2×CH₂CH₃), 6.50 (1H, s, vinylH), 7.2-7.8 (9H, m, vinylH + 8×arylH) and 11.15 (1H, br, NHCH₂).

Coupling of (10b) to 4-(methylamino)butanoic acid

4-(Methylamino)butanoic acid .HCl (31mg, 0.2mmol) was dissolved in methanol (5ml), then a solution of tetra-n-butylammonium hydroxide in methanol (1M, 0.4ml, 0.4mmol) added. To this was added (10b) (60mg, 0.1mmol), giving a brownish solution. T.l.c. (silica; 15% methanol / dichloromethane. (10b) purple, $R_f = 0.4 \rightarrow$ (10d) brown, $R_f < 0.1$). The solvent was then removed under reduced pressure and the residue purified by preparative t.l.c. (silica; methanol 40% ; chloroform 60%) to give the acid dye (10d), 20mg.

λ_{\max} (MeOH) 788nm



(10a) R=O⁻
 (10b) R=OMe MeOSO₃⁻
 (10c) R=NHBU
 (10d) R=N(Me)CH₂CH₂CH₂CO₂H

EXAMPLE 11**Synthesis of phosphoramidite derivatives of squarate dyes****5 Synthesis of dye phosphoramidite (11c)****3-Methoxy-4-(1-ethyl-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)cyclobut-3-en-1,2-dione, sodium salt (11a)****1-Ethyl-2,3,3-trimethylindoleninium-5-sulphonate (1e)**

10 (1.33g, 5mmol) was dissolved in dry methanol (12ml); to the resulting solution was added sodium methoxide (0.27g, 5mmol) and the mixture stirred until all the solid had dissolved (5min). 3,4-Dimethoxycyclobut-3-en-1,2-dione (0.71g, 5mmol) was then added and the resulting mixture heated at reflux under nitrogen atmosphere for 4h. The greenish-yellow
 15 mixture was then cooled at 0°C for 16h. The precipitated yellow solid was collected by filtration, washed with ice-cold ethanol and diethyl ether, then dried under vacuum at 50°C to give the *title compound* (11a), 0.67g (34%).

λ_{\max} (MeOH) 422nm

20 δ_{H} (270MHz, CD₃OD) 1.3 (3H, t, indole N-CH₂-CH₃), 1.6 (6H, s, indole CMe₂), 4.0 (2H, g, indole N-CH₂-CH₃), 4.6 (3H, s, -OMe), 5.6 (1H, s, methine -CH=), 7.1 (1H, d), 7.8 (2H, m).

25 **2-(1-Ethyl-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-4-(1-(5-hydroxypentyl)-3,3-dimethyl-2-indolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (11b)**

Intermediate (1m) (415mg, 1.0mmol) and half-dye (11a) (350mg, 0.89mmol) were mixed in dry 1-butanol (10ml) and the mixture heated at reflux for 6h, giving a deep blue solution. The reaction was then deemed
 30 to be complete by UV/VIS (methanol solution, λ_{\max} 632nm) and t.l.c. (C-

18 silica; ethanol 50% : water 50%. One blue spot at $R_f = 0.3$.

The acetate-dy was not isolated; the solvent was removed under reduced pressure and the residue redissolved in methanol (5ml).

To this solution was added potassium carbonate (200mg) and the mixture stirred for about 1h (C-18 silica; ethanol 50% : water 50%. Dye-OAc, $R_f = 0.3 \rightarrow$ (11b), $R_f = 0.4$). This mixture was purified by prep. HPLC (C-18 column; water \rightarrow methanol gradient) to give the *title compound* (11b), 450mg.

λ_{\max} (MeOH) 632nm

δ_H (300MHz, DMSO-broad spectrum) 1.26 (3H, t, indole N-CH₂-CH₃), 1.45 (4H, m, 2x -CH₂-), 1.67 (14H, 2x indole CMe₂ and -CH₂-), 3.38 (2H, -CH₂-OH), 4.09 (4H, 2x indole N-CH₂-), 4.39 (1H, -OH), 5.77+5.80 (each 1H, s, 2x methine -CH=), 7.16 (1H, m) + 7.24 (1H, d) + 7.33 (2H, m) = 4x indole aryl-H, 7.51 (1H, d) + 7.59 (1H, d) + 7.66 (1H, s) = 3x sulphonated indole aryl-H.

Phosphitylation to give phosphoramidite dye (11c)

The hydroxy dye (11b) (250mg) was dissolved in dry N,N-dimethylformamide (5ml). To the resulting solution was added N,N-diisopropylethylamine (0.17ml) and 2-cyanoethyl-N,N-diisopropylchlorophosphoramidite, in 0.05ml aliquots with t.l.c. monitoring (C-18 silica; ethanol 50% : water 50%. (11b), $R_f = 0.4 \rightarrow$ (11c), $R_f = 0.5$). After addition of three aliquots (0.15ml total) the reaction appeared complete. This solution was then used directly on an automated DNA synthesiser.

Synthesis of dye phosphoramidite (11f)**2-(1-Ethyl-3,3-dimethyl-2-indolinyliidenemethyl)-4-(1-(5-acetoxypentyl)-3,3-dimethyl-2-indolinyliidenemethyl)-cyclobutenediylum-1,3-diolate**5 **(11d)**

Intermediate (1m) (415mg, 1.0mmol) and 3-butoxy-4-(1-ethyl-3,3-dimethyl-2-indolinyliidenemethyl)-cyclobut-3-en-1,2-dione (7b) (339mg, 1.0mmol) were mixed with anhydrous 1-butanol (10ml); the resulting mixture was then heated at reflux for 16h, giving a deep blue solution. The solvent was removed under reduced pressure and the residue purified by flash chromatography (silica; 4-10% methanol / dichloromethane) to give the *title dye* (11d), 420mg.

 λ_{\max} (MeOH) 628nm

δ_{H} (300MHz, CDCl_3) 1.38 (3H, t, J 7.1Hz, indole N- CH_2 - CH_3), 1.47 (2H, m, $-\text{CH}_2-$), 1.65-1.85 (16H, m, $2\times$ indole CMe_2 and $2\times$ $-\text{CH}_2-$), 2.02 (3H, s, CH_3 -COO-), 4.00 (4H, broad, $2\times$ indole N- CH_2-), 4.04 (2H, t, J 6.4Hz, $-\text{CH}_2$ -OAc), 5.92-5.94 (each 1H, s, $2\times$ methine $-\text{CH}=\text{CH}-$), 6.96 (2H, m) + 7.13 (2H, m) + 7.30 (4H, m) = $8\times$ indole aryl-H.

20 **2-(1-Ethyl-3,3-dimethyl-2-indolinyliidenemethyl)-4-(1-(5-hydroxypentyl)-3,3-dimethyl-2-indolinyliidenemethyl)-cyclobutenediylum-1,3-diolate**
(11e)

To a solution of acetate-protected dye (11d) (400mg) in methanol (20ml) was added potassium carbonate (200mg). The mixture was stirred at room temperature with t.l.c. monitoring (silica; 10% methanol / dichloromethane. (11d), $R_f = 0.6 \rightarrow$ (11e), $R_f = 0.35$). After 2h the solvent was removed under reduced pressure and the residue was purified by flash chromatography (silica; 4-10% methanol / dichloromethane) to give the *title compound* (11e), 338mg.

λ_{max} (MeOH) 628nm

δ_{H} (300MHz, CDCl_3) 1.37 (3H, t, J 7.2Hz, indole $\text{N-CH}_2\text{-CH}_3$), 1.54 (2H, m, $-\text{CH}_2-$), 1.67-1.88 (16H, m, $2 \times$ indole CMe_2 and $2 \times -\text{CH}_2-$), 2.45 (1H, broad s, $-\text{OH}$), 3.67 (2H, t, J 6.2Hz, $-\text{CH}_2\text{-OH}$), 4.04 (4H, broad, $2 \times$ indole N-CH_2-), 5.92+5.98 (each 1H, s, $2 \times$ methine $-\text{CH}=\text{}$), 6.97 (2H, m) + 7.12 (2H, m) + 7.30 (4H, m) = $8 \times$ indole aryl-H.

Phosphitylation to give phosphoramidite dye (11f)

The hydroxy derivatised dye (11e) (205mg, 0.5mmol) was dissolved in dry dichloromethane (5ml) under nitrogen atmosphere. To the blue solution was added N,N -diisopropylethylamine (0.1ml), followed by 2-cyanoethyl- N,N -diisopropylchlorophosphoramidite (5 drops), and the mixture stirred at room temperature for 1.5h. T.l.c. (silica; 10% methanol / dichloromethane. (11e), $R_f = 0.35 \rightarrow$ (11f), $R_f = 0.55$). The mixture was then diluted with 10% triethylamine / ethyl acetate and washed with 10% aqueous sodium carbonate solution and brine. The organic solution was dried (MgSO_4), filtered and the solvent removed under reduced pressure. The resulting gum was purified through a short silica plug (5-15% triethylamine / dichloromethane) to give the *title compound* (11f) (150mg), after removal of solvent. This solution was then used directly on an automated DNA synthesiser.

Synthesis of phosphoramidite dye (11i)

2-4-Bis-(1-(5-hydroxypentyl)-3,3-dimethyl-2-indolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (11g)

Intermediate (1m) (4.35g, 10.5mmol) and 3,4-dihydroxycyclobut-3-en-1,2-dione (570mg, 5.0mmol) were mixed in dry 1-butanol (20ml); the resulting mixture was heated at reflux for 16h, giving a

deep blue solution. T.l.c. analysis showed three blue products (silica; 15% methanol / dichloromethane. $R_f = 0.8, 0.65$ and 0.55). The solvent was removed under reduced pressure and the residue dissolved in methanol (20ml), then potassium carbonate (500mg) added. This mixture was stirred at room temperature for 2h; (silica; 15% methanol / dichloromethane. $R_f = 0.8, 0.65$ and $0.55 \rightarrow R_f 0.55$ only). After removal of solvent the residue was partitioned between water and dichloromethane; the organic layer was retained, washed with water and brine, then dried (Na_2SO_4), filtered and the solvent removed under reduced pressure. The crude dye was purified by flash chromatography (silica; 4-15% methanol / dichloromethane) to give the *title compound* (**11g**), 1.47g.

λ_{max} (MeOH) 630nm

δ_{H} (300MHz, CDCl_3) 1.52 (4H, m, $2 \times -\text{CH}_2-$), 1.65-1.90 (20H, m, $2 \times$ indole CMe_2 and $4 \times -\text{CH}_2-$), 2.7 (2H, broad, $2 \times -\text{OH}$), 3.66 (4H, t, $J 6.2\text{Hz}$, $2 \times -\text{CH}_2-\text{OH}$), 4.02 (4H, broad, $2 \times$ indole $\text{N}-\text{CH}_2-$), 5.96 (2H, s, $2 \times$ methine $-\text{CH}=\text{}$), 6.97 (2H, d, $J 7.7\text{Hz}$) + 7.12 (2H, t, $J 7.3\text{Hz}$) + 7.30 (4H, m) = $8 \times$ indole aryl-H.

2-(5-Hydroxypentyl-3,3-dimethyl-2-indolinyldenemethyl)-4-(1-(4,4'-dimethoxytrityloxy)pentyl)-3,3-dimethyl-2-indolinyldenemethyl)-cyclobutenediylum-1,3-diolate (11h)

The bis-hydroxy dye (**11g**) (1.14g, 2.0mmol) and 4,4'-dimethoxytrityl chloride (0.75g, 2.2mmol) were dissolved in dry pyridine (10ml) and stirred at room temperature for 16h under nitrogen atmosphere. T.l.c. analysis (silica; 10% methanol / dichloromethane) showed three blue spots, corresponding to unreacted (**11g**) ($R_f=0.25$), product (**11h**) ($R_f=0.45$) and some bis-protected compound ($R_f=0.9$). The reaction was quenched by the addition of methanol (1ml) followed by

5mins stirring, before the solvent was removed under reduced pressure. The blue components were then separated by flash chromatography (silica; 1-10% methanol / chloroform). Fractions containing the bis-protected compound were combined and treated with trichloroacetic acid (0.5g) for 1hr, then combined with those containing unreacted dye. (11g) was then recovered and re-reacted as above. After reaction and work-up, the product (11h) was combined with that from the first reaction to give a total of 1.18g.

λ_{max} (CH_2Cl_2) 634nm. Addition of trichloroacetic acid gave extra peaks at 416nm and 504nm, corresponding to DMT cation.
 δ_{H} (300MHz, CDCl_3) 1.02 (4H, m, $2\times -\text{CH}_2-$), 1.51-1.88 (20H, m, $2\times$ indole CMe_2 and $4\times -\text{CH}_2-$), 2.4 (1H, broad, $-\text{OH}$), 3.03 (2H, t, J 6.0Hz, $-\text{CH}_2-\text{ODMT}$), 3.66 (2H, t, J 6.2Hz, $-\text{CH}_2-\text{OH}$), 3.76 (6H, s, $2\times \text{Ar}-\text{OMe}$), 4.00 (4H, broad, $2\times$ indole $\text{N}-\text{CH}_2-$), 5.92+5.98 (each 1H, s, $2\times$ methine $-\text{CH}=\text{}$), 6.78 (4H, m), 6.94 (2H, m), 7.09-7.40 (15H, m).

Phosphitylation to give phosphoramidite dye (11i)

Dye (11h) (220mg, 0.25mmol) was dissolved in dry tetrahydrofuran (2ml); to this deep-blue solution was added N,N-diisopropylethylamine (0.1ml), followed by 2-cyanoethyl-N,N-diisopropylchlorophosphoramidite (0.1ml, 0.4mmol). The resulting mixture was stirred at room temperature and the reaction monitored by t.l.c. (silica; ethyl acetate 50% : acetonitrile 50%. (11h), $R_f = 0.35 \rightarrow$ (11i), $R_f = 0.65$). After 2h the mixture was diluted with ethyl acetate (25ml), washed with 5% aqueous sodium hydrogen carbonate solution and brine, then dried (MgSO_4) and filtered through a 0.5cm-thick plug of silica, washing through with ethyl acetate. The solvent was removed under reduced pressure, then re-evaporated from dry toluene solution. The residue was then dried under high vacuum, re-analyzed by t.l.c. (silica; ethyl acetate

50% : acetonitrile 50%. One blue spot, $R_f = 0.65$) and used directly for DNA labelling on an automated DNA synthesiser.

Synthesis of dye phosphoramidite (11i)

2-4-Bis-(1-(5-hydroxypentyl)-3,3-dimethyl-2-benzindolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (11j)

Synthesised by an analogous method to dye (11g), 2.96g (43%).

λ_{\max} (MeOH) 663nm

δ_H (300MHz, $CDCl_3$) 1.50 (4H, m, 2x $-CH_2-$), 1.66 (4H, m, 2x $-CH_2-$), 1.86 (4H, m, 2x $-CH_2-$), 1.97 (12H, s, 2x indole CMe_2), 3.4 (2H, s, 2x $-OH$), 3.62 (4H, t, 2x $-CH_2-OH$), 4.07 (4H, broad, 2x indole $N-CH_2-$), 5.97 (2H, s, 2x methine $-CH=$), 7.22 (2H, d), 7.32 (2H, app t), 7.47 (2H, td), 7.79 (4H, app t), 8.10 (2H, d).


2-(5-Hydroxypentyl-3,3-dimethyl-2-benzindolinyliidenemethyl)-4-(1-((4,4'-dimethoxytrityloxy)pentyl)-3,3-dimethyl-2-benzindolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (11k)

Synthesised by an analogous method to dye (11h), 1.58g (38%).



λ_{\max} (MeOH) 663nm

MS (MALDI-TOF): 939


δ_H (300MHz, $CDCl_3$) 1.4-1.9 (12H, m, 6x $-CH_2-$), 1.982+1.985 (each 6H, s, 2x indole CMe_2), 2.98 (2H, t, $-CH_2-ODMT$), 3.63 (2H, t, $-CH_2-OH$), 3.67 (3H, s, $MMT-OMe$), 4.04 (4H, broad, 2x indole $N-CH_2-$), 5.93+5.99 (each 1H, s, 2x methine $-CH=$), 6.69 (2H, m), 7.1-7.24 (10H, m), 7.28-7.36 (6H, m), 7.45-7.53 (2H, m), 7.70-7.83 (4H, m), 8.08-8.16 (2H, m).





(11a)



(7b)



(11g) $R_1=R_2=H$ (11h) $R_1=H$ $R_2=ODMT$ (11i) $R_1=P(OCH_2CH_2CN)N^iPr_2$ $R_2=ODMT$ (11j) $R_1=R_2=H$ (11k) $R_1=H$ $R_2=ODMT$ (11l) $R_1=P(OCH_2CH_2CN)N^iPr_2$ $R_2=ODMT$

Incorporation of dye monomers into DNA primers and use in sequencing

- 5 Dye primers were synthesised as follows on an ABI 394 4-column DNA synthesiser using Pharmacia "Pac" base amidites [Example (11f)] or Glen Research "Ultra-Mild" amidites [(11c), (11i)]. All other synthesis reagents were from ABI. Oligonucleotides were prepared on a 0.2 μ mol scale using the standard cycle except for the dye-amidite
- 10 coupling reaction where the coupling time was manually extended.

The primer sequence used was a -21 M13 universal 18mer 5' tgtaaacgacggccagt 3'. The dye phosphoramidites were added to the 5' terminus. Cleavage from the support and subsequent deprotections were performed using either 30% NH_4OH at 60 °C for 20 min (11f) or 5 0.05 M K_2CO_3 in MeOH for 2h at 25 °C. After deprotection the crude oligonucleotides were concentrated under vacuum and then precipitated by addition of 1/10th volume of 3M sodium acetate and 3 volumes of absolute ethanol. After centrifugation at 13000g for minutes the DNA pellets were washed with 70% ethanol and dissolved in 100 μl of 95% TE 10 buffer/5% acetonitrile ready for HPLC. A fraction was used for spectrophotometric analysis - an approximation of percentage labelling was estimated from the ratio of the DNA to dye extinction coefficients. Final purification of the oligonucleotides was performed by HPLC using Spherisorb ODS2 C18 column [5 μ] and 0.1M ammonium acetate and 15 acetonitrile as eluent with a 5-70% acetonitrile gradient at 1 ml/min. Detection was performed at 260 and 640 nm with collection of fractions absorbing at both wavelengths. These were concentrated under vacuum and ethanol precipitated as above. The DNA pellets were dissolved in TE buffer and the OD 215-750 nm spectrum determined. Primers were 20 diluted to 2 pmol/ μl for DNA sequencing.

Sequencing experiments were performed as outlined in Example 5 with excellent results.

EXAMPLE 12

Synthesis of hydroxy squarate dye derivatives suitable for
conversion to phosphoramidites.

2-(1-(13-Hydroxy-7-aza-6-oxotridecanyl)-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-4-(1-ethyl-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (12a)

Activation of dye (2b) as the NHS ester

Dye (2b) (100mg, 0.11mmol), O-(N-succinimidyl)-N,N,N',N'-bis(tetramethylene) uronium hexafluorophosphate (70mg, 0.17mmol) and N,N-diisopropylethylamine (10 drops) were mixed in dry N,N-dimethylformamide (2ml) to give a deep blue solution. The reaction to give the dye-NHS ester was monitored by t.l.c. (C-18 silica; methanol 40% : water 60%. (2b) R_f = 0.2, NHS ester R_f = 0.3). The reaction was complete after 2h. The product was used in the next reaction without any further manipulations.

2-(1-(13-Hydroxy-7-aza-6-oxotridecanyl)-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-4-(1-ethyl-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (12a)

To the above mixture was added 6-aminohexan-1-ol (25mg, 0.2mmol) and the mixture stirred at room temperature with t.l.c. monitoring (C-18 silica; methanol 50% : water 50%. NHS ester R_f = 0.4, (12a) R_f = 0.55). Reaction was complete after 1h. The crude dye was precipitated with diethyl ether, dried under vacuum, then purified by prep. HPLC (C-18 silica column; water→methanol gradient). Yield of the titled compound (12a) = 70mg.

2-(1-(14-(4,4'-Dimethoxytrityloxy)-13-hydroxy-11-oxa-7-aza-6-oxotetradecanyl)-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-4-(1-ethyl-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (12b)

5

Activation of dye (2b) as the NHS ester


Dye (2b) (40mg, 0.05mmol) was reacted as described in Example (12a), to give the NHS ester which was used without purifying.

10 2-(1-(14-(4,4'-Dimethoxytrityloxy)-13-hydroxy-11-oxa-7-aza-6-oxotetradecanyl)-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-4-(1-ethyl-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (12b)


To the above mixture was added the 7-(4,4'-
15 dimethoxytrityloxy)-6-hydroxy-4-oxaheptylamine (30mg, 0.067mmol) and stirring continued. The reaction was monitored by t.l.c. (C-18 silica; ethanol 50% : water 50%. NHS ester R_f = 0.8, (12b) R_f = 0.6). After 1h the reaction was complete and the dye was precipitated by adding diethyl ether. The crude product was purified by prep. t.l.c. (C-18 silica; methanol
20 60% : water 40%) to give pure dye (12b), 55mg.

λ_{max} (MeOH) 636nm; λ_{em} (MeOH)

δ_H (300MHz, DMSO) 1.1-1.35 (5H, m), 1.4-1.7 (18H, m, indole CMe₂ and 3x-CH₂-), 1.97 (2H, t, R-CH₂-CONH-), 2.85 (2H, m, glycol -CH₂-), 2.94 (2H, m, glycol -CH₂-), 3.10 (1H, m, glycol -CH-), 3.28
25 (4H, m, -CONH-CH₂- and -CH₂O-glycol), 3.64 (6H, s, 2xAr-OMe), 4.04 (4H, broad m, 2x indole N-CH₂-), 4.81 (1H, appd, -OH), 5.74 (2H, s, 2x methine -CH=), 6.79 (4H, m) + 7.08-7.28 (9H, m) = 13x DMT aryl-H, 7.32 (2H, m) + 7.55 (2H, m) + 7.61 (2H, m) = 6x indole aryl-H, 7.69 (1H, t, -CONH-).



(2b) R=H

(12a) R=NH(CH₂)₆OH

(12b)

EXAMPLE 13

Synthesis of non sulphonated squarate dyes and comparison of photostability properties with sulphonic acid derivatised squarate dyes.

2-(1-(5-Carboxypentyl)-3,3-dimethyl-2-indolinyldenemethyl)-4-(1-ethyl-3,3-dimethyl-2-indolinyldenemethyl)cyclobutenediylum-1,3-diolate (13a)

The title compound was prepared as its butyl ester in a similar manner to (2b) butyl ester, *vide supra*, (1i), squaric acid and (1g).

λ_{max} (MeOH) 630 nm; λ_{em} (MeOH) 643 nm

δ_{H} (270 MHz; CDCl₃) 0.93 (3H, t, J 7.3), 1.3-1.92 (13H, m), 1.78 (12H, s), 2.32 (2H, t, J 7.3), 3.9-4.2 (4H, m), 4.06 (2H, t, J 6.7), 5.94 (1H, s), 5.97 (1H, s), 6.95-7.05 (2H, m), 7.11-7.25 (2H, m) and 7.26-7.45 (4H, m);

Saponification of the butyl ester in an analogous method to Example (2b) gave the title compound (13a)

δ_{H} (270 MHz; CDCl₃) 1.40 (3H, t, J 7.3), 1.60-2.00 (6H, m), 1.78 (12H, s), 2.42-2.50 (2H, m), 3.95-4.18 (4H, m), 5.90-6.00 (1H, br s), 6.05-6.15 (1H, br s), 6.93-7.08 (2H, m), 7.10-7.25 (2H, m) and 7.28-7.45 (4H, m)

Activation of (13a) to an succinimidyl ester was carried out as per the method described in Example (4)

2-(1-Butyl-3,3-dimethyl-5-carboxymethyl-2-indolinyldenemethyl)-4-(1-ethyl-3,3-dimethyl-2-indolinyldenemethyl)cyclobutenediylum-1,3-diolate (13b)

The title compound was prepared as its butyl ester in a similar manner to Example (2b) butyl ester, *vide supra*, (1j), squaric acid

and (1g).

λ_{max} (MeOH) 632 nm

δ_{H} (270 MHz; CDCl_3) 0.90 (3H, t), 0.98 (3H, t), 1.20-2.10 (7H, m), 1.80 (12H, s), 3.65 (2H, br s), 3.90-4.20 (6H, m), 5.95 (2H, s), 6.85-7.05 (2H, m) and 7.15-7.45 (5H, m)

Saponification of the butyl ester in an analogous method to Example (2b) gave the title compound (13b)

δ_{H} (270 MHz; CDCl_3) 0.95 (3H, t), 1.35-1.55 (5H, m), 1.70-1.90 (14H, m), 3.70 (2H, s), 3.80-4.20 (4H, m), 5.95 and 6.00 (each 1H, s), 6.92 (1H, d), 7.00 (1H, d), 7.15 (1H, d) and 7.32-7.41 (4H, m)

Activation of (13a) to an succinimidyl ester was carried out as per the method described in Example (4)

2-(1-(5-Carboxypentyl)-3,3-dimethyl-2-benzindolinyliidenemethyl)-4-(1-ethyl-3,3-dimethyl-2-benzindolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (13c)

A mixture of (1k) (119 mg), (1l) (162 mg) squaric acid (37 mg) and potassium acetate (98 mg) in 2-butanol (20 ml) was heated at 100 °C for 10 h and then concentrated *in vacuo*. Purification of the residue by HPLC (C_{18} isocratic MeOH) afforded the title compound (13c).

δ_{H} (270 MHz; CDCl_3) 1.45 (3H, t), 1.76 (2H, m), 1.85-2.05 (4H, m), 2.07 (12H, s), 2.48 (2H, m), 4.10-4.28 (4H, m), 5.98 and 6.12 (each 1H, s),

Activation of (13c) to an succinimidyl ester was carried out as per the method described in Example (4).

2-(1-(5-Carboxypentyl)-3,3-dimethyl-2-indolinyliidenemethyl)-4-(3-ethyl-2-benzothiazolinyliidenemethyl)-cyclobutenediylum-1,3-diolate (13d)

Synthesised from the intermediate half-dye, prepared from intermediate (1l) and squaric acid, and intermediate (1s) to give the n-

butyl ester as per Example (2b).

Saponification gave the free acid (13d) and this was converted to the N-succinimidyl ester as per Example (4).

λ_{\max} (MeOH) 638 nm

5 δ_H (270MHz; $CDCl_3$) 1.46 (3H, t, CH_2Me), 1.63-2.07 (6H, m), 1.75 (6H, s, $2 \times Me$), 2.46 (2H, t, CH_2CO_2H), 3.95 (2H, br t, NCH_2CH_2), 4.25 (2H, q, NCH_2CH_3), 5.93 and 6.00 (each 1H, s, $2 \times$ vinyl CH), 6.90 (1H, d, ArH), 7.04-7.54 (6H, series m, ArH) and 7.61 (1H, d, ArH)

10 2-(1-(5-Carboxypentyl)-3,3-dimethyl-2-benzindolinyldenemethyl)-4-(1-methyl-3,3-dimethyl-2-indolinyldenemethyl)-cyclobutenediylum-1,3-diolate (13e)


Synthesised from the intermediate half-dye, prepared from 1,2,3,3-tetramethylindoleninium iodide and squaric acid, and intermediate

15 (1l) to give the n-butyl ester as per Example (2b).


Saponification gave the free acid (13e)

λ_{\max} (MeOH) 644 nm


20 Free acid: δ_H (300 MHz; $CDCl_3$) 1.70-1.80 (12H, m, indole $CMe_2 + 3 \times CH_2$), 2.05 (6H, s, benzindole CMe_2), 2.47 (2H, br t, CH_2CO_2H), 3.57 (3H, br s, indole NCH_3), 4.16 (2H, br t, benzindole NCH_2), 5.88+6.14 (each 1H, s, vinylH), 7.00 (1H, d), 7.13 (1H, m), 7.26-7.36 (3H, m), 7.43 (1H, app t), 7.58 (1H, app t), 7.89 (2H, app t), 8.20 (1H, d).




(13a) X=CMe₂
(13d) X=S



(13b)



(13c)



(13e)

Photostability of Indolinium Based Squarate Dyes

EXAMPLE	OVERALL CHARGE	NO. OF ARYL SULPHONATE GROUPS	NO. OF ALKYL SULPHONATE GROUPS	T _{1/2} (MIN)
Cy5™	-1	2	0	26.4
13a	0	0	0	25.4
13b	0	0	0	26.1
2f	-1	1	0	40.4
2d	-1	1	0	51.0
2e	-3	1	2	61.2
2c	-2	2	0	81.3
2a	-3	2	1	88.5
2b	-2	2	0	94.9

5

Benzindole Based Dyes

EXAMPLE	OVERALL CHARGE	NO. OF ARYL SULPHONATE GROUPS	NO. OF ALKYL SULPHONATE GROUPS	T _{1/2} (MIN)
13c	0	0	0	22.4
2h	-2	2	0	38.9

Photostability of Mixed Thiazolinium/Indolinium Based Squarate**Dyes**

EXAMPLE	OVERALL CHARGE	NO. OF ARYL SULPHONATE GROUPS	NO. OF ALKYL SULPHONATE GROUPS	T _{1/2} (MIN)
13d	0	0	0	7.7
3b	-1	0	1	10.7
3a	-1	1	0	15.2

5

EXAMPLE 14**Synthesis of an Energy Transfer Cassette**

10 **2,4-bis(1-(5-Carboxypentyl)-3,3-dimethyl-5-sulphonato-2-indolinyliidenemethyl)cyclobutenediylum-1,3-diolate (14a)**

Intermediate (1c) (1.42g, 3.0mmol) and squaric acid (0.17g, 1.5mmol) were mixed in anhydrous 1-butanol (10ml). The resulting mixture was heated at 110°C for 65h. A deep green-blue colour was
 15 generated during this time, with some dark solid present. The solvent was removed under reduced pressure; the residue was then redissolved in methanol (15ml), and a solution of potassium hydroxide (0.5g, 9.0mmol) in water (10ml) added. This solution was stirred for 20h. T.l.c. analysis (C18-silica; methanol 50%: water 50%. Major blue spot at R_f=0.6). The
 20 solution was then neutralised with acetic acid before purification by prep. HPLC (C18, water→methanol gradient) to give *title dye* (14a).

λ_{max} (MeOH) = 638nm; λ_{ex} (MeOH) = 636nm; λ_{em} = 644nm.

δ_{H} (300MHz, CD₃OD) 1.52 (4H, m), 1.70 (4H, m), 1.77 (12H, s, indole CMe₂), 1.86 (4H, m), 2.35 (4H, t, J 7.3, 2x -CH₂CO₂H), 4.17 (4H, broad s, 2x indole N-CH₂-), 6.03 (2H, s, 2x vinyl -CH=), 7.33 (2H, d, J 7.7Hz), 7.86-7.90 (4H, m).

25

3-(3-(t-Butyloxycarbonylamino)propylamino)-2,4-bis(3-ethyl-2-benzothiazolinyliidenemethyl)-cyclobutenediylum-1-olate methosulphate (14b)

Dye (9b) (560mg, 1.0mmol) was dissolved in
5 dichloromethane (20ml). To the resulting deep blue solution was added t-butyl-N-(3-aminopropyl) carbamate (190mg, 1.1mmol). This mixture was stirred at room temperature until the reaction was complete by t.l.c. (silica; 10% methanol in dichloromethane. (9b) $R_f=0.2 \rightarrow$ (14b) $R_f=0.3$).
The solution was then washed with water, dried (Na_2SO_4), filtered and the
10 solvent removed under reduced pressure to a volume of about 5ml. The product was then precipitated by addition of diethyl ether; the resulting solid was collected, washed with ether and dried under vacuum at 35°C to give the *title dye* (14b) (640mg).

λ_{max} (MeOH) = 658nm; λ_{ex} (MeOH) = 658nm; λ_{em} = 670nm.
15 δ_{H} (300MHz, CDCl_3) 1.40 (9H, s, -CMe₃), 1.48 (6H, t, J 7.0Hz, thiazole N-CH₂-CH₃), 1.97 (2H, quin, -CH₂-CH₂-CH₂-), 3.31 and 3.66 (each 2H, q, NH-CH₂-), 3.79 (3H, s, methosulphate), 4.13 and 4.44 (each 2H, q, thiazole N-CH₂-CH₃), 5.69 (1H, s, vinyl-CH=), 5.89 (1H, broad t, BOC-NH-), 6.58 (1H, s, vinyl-CH=), 7.12-7.20 (2H, m), 7.28-7.36
20 (3H, m), 7.42-7.47 (2H, m), 7.58 (1H, d), 8.83 (1H, broad t, squarate-NH-CH₂-).

3-((Aminopropyl)amino)-2,4-bis(3-ethyl-2-benzothiazolinyliidenemethyl)-cyclobutenediylum-1-olate trifluoroacetate, trifluoroacetic acid salt (14c)

The protected dye (14b) (250mg) was dissolved in
chloroform (4ml). To the deep blue solution was added trifluoroacetic acid
(2ml), turning the solution a yellow-brown colour. This mixture was stirred
for 1h, the solvent was removed under reduced pressure. The residue
30 was redissolved in 4ml of 10% methanol / dichloromethane, restoring the

blue colour. The dy was precipitated by addition of diethyl ether; the solid was collected, washed well with fresh ether and dried under vacuum at 50°C to give the *title amine dye* (14c), 245mg.

λ_{max} (MeOH) = 656nm; λ_{ex} (MeOH) = 657nm; λ_{em} = 671nm.

5 δ_{H} (300MHz, DMSO) 1.27-1.36 (6H, 2x overlapping t, thiazole N-CH₂-CH₃), 1.95 (2H, quin, -CH₂-CH₂-CH₂-), 2.96 (2H, broad, -CH₂-NH₃⁺), 3.72 (2H, q, -NH-CH₂-), 4.33-4.45 (4H, , 2x overlapping broad q, thiazole N-CH₂-CH₃), 5.91 and 6.23 (each 1H, s, vinyl-CH=), 7.32-7.42 (2H, app quin.), 7.49-7.58 (2H, m), 7.65-7.75 (2H, 2xd), 7.84 (3H, broad s, -NH₃⁺), 7.94-8.02 (2H, 2xd), 8.98 (1H, broad t, squarate-NH-CH₂-).

Coupling of (14a) to (14c) to give the ET cassette (14d)


Diacid dye (14a) (60mg) was dissolved in anhydrous N,N-dimethylformamide (1ml); to the deep blue solution was added O-(N-succinimidyl)-N,N,N',N'-bis(tetramethylene)uronium hexafluorophosphate
15 (27mg) and N,N-diisopropylethylamine (50μl). This mixture was stirred at room temperature for 1h. T.l.c. (C18 silica; methanol 50% : water 50%. (14a) R_f=0.5→mono-NHS ester, R_f=0.35 and bis-NHS ester, R_f=0.2). To the above mixture was then added N,N-diisopropylethylamine (150μl) and
20 amine dye (14c) (45mg); this mixture was stirred for another 2h. T.l.c. (C18 silica; methanol 85% : water 15%. (14a) R_f=0.95, amine (14c) R_f=0.1, product (14d) R_f=0.55). The crude dye was isolated by precipitation with diethyl ether. The crude solid was purified by prep. HPLC (C18 silica; water→methanol gradient) to give the *compound dye*
25 (14d).


λ_{max} (MeOH) = 638nm with shoulders to short and long wavelengths.

λ_{ex} (MeOH) = 640nm; λ_{em} = 671nm.

30 δ_{H} (300MHz, DMSO) 1.17-1.35 (8H, m, 2x thiazole N-CH₂-CH₃ and -CH₂-), 1.50-1.72 (24H, m, 2x indole N-CH₂-CH₃ and 6x -CH₂-).


2.03 (2H, broad t, $-\text{CH}_2-\text{CONH}$), 2.18 (2H, t, 7.2Hz, $-\text{CH}_2-\text{CO}_2\text{H}$), 3.07 and 3.52 (each 2H, broad, $\text{NH}-\text{CH}_2-$), 4.04 (4H, broad, $2\times \text{indole N}-\text{CH}_2-\text{CH}_3$), 4.38 (4H, broad, $2\times \text{thiazole N}-\text{CH}_2-\text{CH}_3$), 5.77 (2H, s, indole vinyl-H), 5.85 and 6.31 (each 1H, s, thiazole vinyl-H), 7.21-7.39 (4H, m, $4\times \text{aryl-H}$), 7.43-7.72 (8H, m, $8\times \text{aryl-H}$), 7.87-7.99 (3H, m, $2\times \text{aryl-H}$ and $-\text{CONH}-$), 8.82 (1H, broad t, squarate $-\text{NH}-\text{CH}_2-$).





(14b) $R = \text{NH-CO.O}(\text{CH}_3)_3$

(14c) $R = \text{NH}_3^+ \text{CF}_3\text{CO}_2^-$



EXAMPLE 15**Synthesis of a Squarate Dye Labelled Peptides Using Both Solution and Solid Phase Approaches****Representative example:****SYNTHESIS OF PEPTIDE ON SOLID PHASE**

The serine protein kinase substrate peptide (NH₂-ARRVTSAAARRS-OH) was synthesised using solid phase Fmoc chemistry, the N-terminal Fmoc group was removed at the end of the synthesis. The peptide was cleaved from 100mg of resin and deprotected using a mixture of trifluoroacetic acid, water, thioanisole and ethanedithiol (95:2.5:5:2.5 v/v, 2 ml) for 90 minutes. The crude peptide was precipitated from cold diethyl ether, centrifuged down, dried in vacuo, then after dissolving in water, purified by semi-preparative HPLC using a Vydac C-18 reverse phase column at a flow rate of 4 ml/minute and a gradient of water/0.1% TFA to 60% acetonitrile/0.1% TFA over a period of 30 minutes. Detection was at 230 nm. A major peak eluting at 8.5 minutes was collected and freeze dried to give 10 mg of the desired peptide as a white solid.

PREPARATION OF DYE-NHS ESTER

Squarate dye (2i) (88mg, 0.117 mmol), N-hydroxysuccinimide (20mg, 0.174 mmol) and N-cyclohexyl-N'-(2-morpholinoethyl)-carbodiimide methyl-p-toluenesulphonate (45 mg, 0.11 mmol) and a single small crystal of 4-dimethylaminopyridine were placed in a round bottomed flask fitted with a magnetic stirrer bar and dry DMF (2 ml) was added. The mixture was stirred for 16 hours at ambient temperature, then the solvent was removed under vacuum and the blue residue was dissolved in dry DMSO (1 ml).

SOLID PHASE LABELLING




100 mg of peptide resin (equivalent to ~0.03 mmol of peptide) was weighed into a 1.5 ml polypropylene V-vial then 0.4 ml of the dye-NHS ester solution was added followed by 0.6 ml of dry DMSO plus
5 0.02 ml diisopropylethylamine. The vial was placed on rollers in the dark at ambient temperature for 18hrs. The resin was filtered off, washed with 2x1 ml DMSO, 2x1 ml methanol and finally 2x1 ml dichloromethane, then dried *in vacuo*. The resin was treated with 2 ml of the deprotection mixture as outlined above to cleave the labelled peptide from the resin
10 and remove the protecting groups. The peptide was precipitated from diethyl ether as a blue solid. This was treated in the same way as the unlabelled peptide described above. Upon HPLC purification, a blue coloured peak eluted after 22.5 minutes, this was collected and freeze dried to give 3 mg of blue solid. Mass spec gave a peak at 1933 m.u.
15 (calculated mol. wt. of the squarate dye labelled peptide =1930).

SOLUTION PHASE LABELLING

0.3 ml of the dye-NHS solution was added to 5.0 mg (0.004 mmol) of the peptide in a polypropylene V-vial, a further 0.7 ml of dry
20 DMSO plus 0.02 ml of diisopropylethylamine was added, then the vial was placed on rollers in the dark for 18 hours. The mixture was then separated on semi-prep HPLC using the same conditions as outlined above

CLAIMS

1. A squarate dye of the formula (I) or (II) or (IIa)



where each Z is independently O or S or CR¹,₂,

n = 1 - 3,

each of s and m is 0, 1 or 2,

R¹ is lower alkyl (1 - 4 carbon chain),

5 each R² is independently selected from electron donating and electron withdrawing groups or is a branched or straight chain of up to 30 carbon atoms incorporating one to five positively charged nitrogen atoms,




each R³ is independently selected from: alkylene,
10 alkenylene and alkynylene (1 - 20 carbon chain), or is a branched or straight chain of up to 30 carbon atoms incorporating one to five ether oxygen atoms or arylene rings or positively charged nitrogen atoms,

at least one X is a nucleophilic functional group, such as OH, SH or NH₂, or alternatively a grouping capable of reacting with a
15 nucleophile,

and any other X present is independently selected from H and SO₃⁻ and the residue of a squarate dye of formula (I) or (II) or (IIa) and another fluorochrome,

provided that at least one R² is SO₃⁻ and/or at least one X is
20 phosphoramidite.

2. A squarate dye of the formula (III) or (IV) or (IVa)



where each Z is independently O or S or CR¹₂,

n = 1 - 3,

each of s and m is 0, 1 or 2,

R¹ is lower alkyl (1 - 4 carbon chain);

5 each R² is independently selected from electron donating and electron withdrawing groups or is a branched or straight chain of up to 30 carbon atoms incorporating one to five positively charged nitrogen atoms,

each R³ is independently selected from: alkylene,
10 alkenylene and alkynylene (1 - 20 carbon chain), or is a branched or straight chain of up to 30 carbon atoms incorporating one to five ether oxygen atoms or arylene rings or positively charged nitrogen atoms,

at least one X is a nucleophilic functional group, such as OH, SH or NH₂, or alternatively a grouping capable of reacting with a
15 nucleophile

and any other X present is independently selected from H and SO₃⁻ and the residue of a squarate dye of formula (III), (IV) or (IVa) and another fluorochrome,

A is O, NR⁴ or S,


20 R⁴ is alkyl, alkenyl, alkynyl or H, and
each Y is independently X or H.

3. A squarate dye according to claim 1 or claim 2, wherein at least one R² is SO₃⁻.

4. A squarate dye as claimed in any one of claims 1 to 3,
25 wherein n is 1 and Z is -C(CH₃)₂.

5. A squarate dye as claimed in any one of claims 1 to 4, wherein 1 to 5 SO₃⁻ groups are present.

6. A squarate dye as claimed in any one of claims 1 to 5, wherein at least one X is selected from CO₂H, activated carboxyl, CO active ester, NCS, O phosphoramidite, NCOCH₂I and



7. A squarate dye as claimed in any one of claims 1 to 6, wherein each R² is individually selected from halogen, alkoxy, primary secondary and tertiary amine, nitro, SO₃⁻ and -R³X.

10 8. An adduct of a biological molecule with a squarate dye according to any one of claims 1 to 7.

9. An adduct of a nucleoside or nucleotide or analogue or oligonucleotide or nucleic acid with a squarate dye of the formula (I) or (II) or (IIa) or (III) or (IV) or (IVa)

15 where each Z is independently O or S or CR¹₂,


n = 1 - 3,

R¹ is lower alkyl (1 - 4 carbon chain),

each R² is independently selected from electron donating and electron withdrawing groups such as halogen, alkoxy, primary
20 secondary and tertiary amino, nitro, SO₃⁻, and -R³-X, or is a branched or straight chain of up to 30 carbon atoms incorporating one to five positively charged nitrogen atoms,

each R³ is independently selected from: alkylene, alkenylene and alkynylene (1 - 20 carbon chain), or is a branched or
25 straight chain of up to 30 carbon atoms incorporating one to five ether oxygen atoms or arylene rings or positively charged nitrogen atoms, at least one X is a nucleophilic functional group, such as

OH, SH or NH₂, or alternatively a grouping capable of reacting with a nucleophile, in which case X is preferably selected from the following CO₂H, activated carboxyl such as acid halide or anhydride, CO active ester, NCS, O phosphoramidite, NC(O)CH₂I and



any other X present is independently selected from H and SO₃⁻ and the residue of a squarate dye (whereby dimers and oligomers of the dyes shown as monomers of formula (I), (II), (IIa), (III), (IV) and (IVa) are envisaged), or other fluorochrome,

each of s and m is 0, 1 or 2,

A is O, NR⁴ or S,

R⁴ is alkyl, alkenyl, alkynyl or H, and

each Y is independently X or H.

10. An adduct as claimed in claim 9, wherein the adduct has the formula Q-N-CO-Sq, where Q is a nucleotide or nucleotide analogue or oligonucleotide residue, and Sq is a residue of a squarate dye, the two being joined by an amide linkage formed between an amine group of Q and a carboxylate group of Sq.

11. An improved fluorescent sequencing method, which comprises using an adduct according to claim 9 or claim 10.

12. A fluorescent labelling complex comprising:

- a first or donor fluorochrome having first absorption and emission spectra;
- a second or acceptor fluorochrome having second absorption and emission spectra, the wavelength of the emission

maximum of said second fluorochrome being longer than the wavelength of the emission maximum of said first fluorochrome, and a portion of the absorption spectrum of said second fluorochrome overlapping a portion of the emission spectrum of said first fluorochrome;

- 5 • at least one linker for covalently attaching said first and second fluorochromes for transfer of resonance energy transfer between said first and second fluorochromes;
- a target bonding group capable of forming a covalent bond with a target compound;
- 10 wherein at least one of the said first and second fluorochromes is a squarate dye as defined in any one of claims 1 to 7.

INTERNATIONAL SEARCH REPORT

Int. Appl. No.
PCT/GB 97/01105A. CLASSIFICATION OF SUBJECT MATTER
IPC 6 C09B57/00 C09B23/00 G01N33/58

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
IPC 6 C09B G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 4 830 786 A (PEASE JOHN ET AL) 16 May 1989 cited in the application see column 2, line 30 - column 3, line 36 see column 5, line 45 - column 6, line 42 ---	1-13
Y	ANALYTICAL CHEMISTRY, vol. 67, 1995, pages 1742-1748, XP002012845 A.J.G. MANCK ET AL: "visible diode laser-induced fluorescence detection in liquide chromatography after precolumn derivatization of amines" cited in the application see page 1743; figure 1 --- -/-	1-13

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- "A" document defining the general state of the art which is not considered to be of particular relevance
"E" earlier document but published on or after the international filing date
"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
"O" document referring to an oral disclosure, use, exhibition or other means
"P" document published prior to the international filing date but later than the priority date claimed

- "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
"&" document member of the same patent family

Novelty
Obvious

Date of the actual completion of the international search

14 August 1997

Date of mailing of the international search report

26.08.97

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Dauksch, H

Form PCT/ISA/210 (second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Int. l. Application No
PCT/GB 97/01105

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP 0 214 847 A (SYNTEX INC) 18 March 1987 cited in the application see page 13, line 2 - line 32 ---	1-13
A	DE 39 12 046 A (UNIV CARNEGIE MELLON) 15 March 1990 see abstract see page 9, line 3 - page 12, line 65 & US 5 268 486 A cited in the application ---	1-13
A	WO 93 09956 A (POLAROID CORP) 27 May 1993 see abstract ---	3,4
A	WO 93 09172 A (EASTMAN KODAK CO) 13 May 1993 cited in the application see abstract see page 50 - page 56; examples 23-64 -----	1-13

Form PCT/ISA/210 (continuation of second sheet) (July 1992)

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/GB 97/01105

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 4830786 A	16-05-89	US 5416214 A	16-05-95
		US 5039818 A	13-08-91
		US 5310922 A	10-05-94
		US 5329019 A	12-07-94
EP 0214847 A	18-03-87	US 4806488 A	21-02-89
		AU 605109 B	10-01-91
		AU 6236586 A	12-03-87
		CA 1304076 A	23-06-92
		DE 3684741 A	14-05-92
		JP 7037984 B	26-04-95
		JP 62103572 A	14-05-87
DE 3912046 A	15-03-90	JP 2191674 A	27-07-90
		US 5486616 A	23-01-96
		US 5569766 A	29-10-96
		US 5569587 A	29-10-96
		US 5268486 A	07-12-93
WO 9309956 A	27-05-93	US 5227498 A	13-07-93
		US 5227499 A	13-07-93
		CA 2123991 A	27-05-93
		DE 69213020 D	26-09-96
		DE 69213020 T	13-02-97
		EP 0613422 A	07-09-94
		JP 7501497 T	16-02-95
		US 5492795 A	20-02-96
		US 5354873 A	11-10-94
WO 9309172 A	13-05-93	AT 146503 T	15-01-97
		CA 2121507 A	13-05-93
		DE 69216114 D	30-01-97
		DE 69216114 T	10-04-97
		EP 0616621 A	28-09-94
		EP 0700961 A	13-03-96
		ES 2095494 T	16-02-97
		JP 7507074 T	03-08-95
		US 5461136 A	24-10-95
		US 5553714 A	10-09-96